

## Pattern Recognition System for Automating Medical Diagnosis Based on Image Data

Evi Irianti<sup>1</sup>, Nina Anis<sup>2</sup>, and Saifiullah Aziz<sup>3</sup>

<sup>1</sup> Poltekkes Kemenkes Medan, Indonesia

<sup>2</sup> Monash University, Malaysia

<sup>3</sup> Herat University, Afghanistan

---

### Corresponding Author:

Evi Irianti,

Jl. Jamin Ginting KM. 13.5 Laucih Subdistrict, Medan Tuntungan District, Medan, North Sumatra Province, Indonesia.

Email: evidesman@gmail.com

### Article Info

Received: Okt 22, 2024

Revised: Dec 23, 2024

Accepted: Jan 28, 2025

Online Version: April 28, 2025

### Abstract

The increasing volume and complexity of medical image data have presented significant challenges for healthcare professionals in delivering timely and accurate diagnoses. Traditional diagnostic processes are often time-consuming and prone to human error, underscoring the need for automated solutions. This study aims to develop a pattern recognition system to automate medical diagnosis using image data, thereby improving diagnostic accuracy and efficiency. A hybrid methodology was employed, combining image preprocessing, feature extraction using convolutional neural networks (CNNs), and classification through deep learning algorithms. The system was trained and validated using publicly available medical image datasets across various disease types. The results demonstrate high diagnostic accuracy, with the system achieving over 92% precision in identifying disease patterns from image inputs. Furthermore, the model exhibited robustness across different imaging modalities, such as X-rays, MRIs, and CT scans. These findings suggest that the proposed pattern recognition system can serve as a reliable support tool for medical practitioners. In conclusion, the integration of image-based pattern recognition in medical diagnostics holds significant promise in enhancing clinical decision-making processes and reducing diagnostic errors.

**Keywords:** pattern recognition, medical diagnosis, image data, deep learning, convolutional neural networks



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

How to cite: Irianti, E., Anis, N., & Aziz, S. (2025). Pattern Recognition System for Automating Medical Diagnosis Based on Image Data. *Sciencetchno: Journal of Science and Technology*, 4(1), 40-47. <https://doi.org/10.70177/Sciencetchno.v4i1.2126>

Published by: Yayasan Pendidikan Islam Daarut Thufulah

---

## INTRODUCTION

Medical diagnosis has undergone a significant transformation in recent decades due to the rapid advancement of imaging technologies and data processing methods. The increasing availability of medical image data, such as X-rays, MRIs, and CT scans, has paved the way for integrating computational systems into diagnostic practices. The complexity and volume of this image data have made manual interpretation both time-consuming and prone to human error.

Health care systems worldwide are facing growing demands for accurate, fast, and scalable diagnostic solutions. Conventional diagnostic approaches often rely heavily on the expertise of medical professionals, which can lead to variability in outcomes and delays in decision-making (De Meyer et al., 2025; Shashar et al., 2025; Williams et al., 2025). The incorporation of computer-based systems that utilize artificial intelligence and pattern recognition techniques offers the potential to address these challenges.

Pattern recognition, as a subfield of machine learning, provides the capability to analyze complex visual information and identify patterns that may not be visible to the human eye. This technology has been increasingly applied in various domains, including facial recognition, handwriting analysis, and security systems. Its extension to the field of medical diagnosis represents a promising evolution in healthcare innovation.

The reliance on manual image interpretation in clinical settings introduces inconsistencies that can affect diagnostic reliability and patient outcomes. Human interpretation is often influenced by subjective judgment, fatigue, and varying levels of expertise among practitioners (Karume et al., 2025; Nieser et al., 2025; Rahmati et al., 2025). These factors highlight the need for standardized, objective tools that can support diagnostic procedures based on image data.

The sheer volume of medical images generated daily surpasses the processing capacity of medical professionals, creating backlogs and delaying critical decisions. Many healthcare systems lack the technological infrastructure to efficiently manage and interpret this growing repository of diagnostic images. This situation underscores the urgent need for automation to alleviate the burden on clinical personnel and improve service delivery.

Despite advancements in artificial intelligence, the integration of pattern recognition systems into practical diagnostic workflows remains limited and fragmented (Bensoussan et al., 2025; Saggarr et al., 2025; Saniee et al., 2025; Stocklassa et al., 2025; Thorpe et al., 2025). The absence of comprehensive frameworks capable of automatically processing and classifying medical image data impedes the realization of a fully automated diagnostic support system. This gap necessitates further exploration and development of effective pattern recognition solutions tailored for medical applications.

This study aims to design and develop a pattern recognition system that can automate medical diagnosis based on image data. The system is expected to identify relevant patterns and classify medical images with high accuracy and reliability. The research focuses on creating a model that is adaptable to various imaging modalities commonly used in healthcare.

Another objective is to evaluate the performance of the proposed system using real-world medical image datasets. The study will measure accuracy, sensitivity, specificity, and processing time as key performance indicators. Comparative analysis with existing diagnostic tools will also be conducted to validate the efficiency and effectiveness of the system.

The final objective is to explore the practical integration of the pattern recognition system into clinical workflows. This includes assessing user-friendliness, scalability, and compatibility with existing healthcare information systems. The study intends to provide a foundation for future applications of AI-based diagnostic tools in routine medical practice.

Existing literature reveals numerous studies on the use of artificial intelligence in medical imaging, yet many focus narrowly on specific diseases or imaging types (Rajalakshmi et al., 2025). Few approaches provide a generalized framework that can accommodate a broad spectrum of diagnostic

tasks across diverse image formats. The lack of unified systems limits the applicability of current models in real-world healthcare settings.

Several previous models demonstrate promising accuracy in controlled environments, but their performance tends to degrade when applied to diverse, noisy, or incomplete real-life datasets. Furthermore, many models are developed with limited interpretability, making it difficult for clinicians to understand or trust the output. This disconnect between algorithmic results and clinical usability hinders adoption in medical contexts.

A critical gap also exists in the integration of pattern recognition systems with user-centered design principles. Most research emphasizes technical performance without considering the practical needs of end-users, such as radiologists or general practitioners. Addressing this gap is essential to ensure that technological advancements translate into meaningful clinical improvements.

This research introduces a novel approach by integrating advanced pattern recognition techniques with real-time medical image classification capabilities. Unlike prior models, the proposed system is designed to be flexible across multiple diagnostic domains, offering a more comprehensive solution for medical image analysis. The model also incorporates explainable AI features to enhance user trust and interpretability.

The study contributes significantly to the body of knowledge by bridging the gap between algorithm development and clinical application. It introduces a practical framework that emphasizes both diagnostic accuracy and system usability. The research also considers interoperability with existing medical records and imaging systems, ensuring smoother implementation in healthcare environments.

The significance of this study lies in its potential to transform diagnostic practices by reducing human error, expediting decision-making, and improving patient outcomes. The integration of pattern recognition into automated diagnostic tools represents a pivotal advancement in medical technology, with wide-ranging implications for clinical efficiency, patient care, and future research directions.

## RESEARCH METHOD

### *Research Design*

This study employed a quantitative research design with an experimental approach to evaluate the effectiveness of a pattern recognition system in automating medical diagnoses based on image data (Peninah et al., 2025; Ranjan et al., 2025). The design was structured to assess system performance using a dataset of medical images representing various diagnostic categories.

### *Research Target/Subject*

The population consisted of publicly available medical image datasets, particularly those related to radiology, dermatology, and pathology. From these datasets, a stratified random sampling method was used to select a total of 1,200 images, ensuring balanced representation across multiple disease categories to maintain data integrity and avoid class imbalance.

### *Research Procedure*

The procedure began with preprocessing the image data, including normalization, resizing, and augmentation to improve training efficiency and model generalization. The dataset was split into training (70%), validation (15%), and testing (15%) sets. The CNN model was then trained using the training set and tuned with the validation set. After training, the model was evaluated using the test set to determine its diagnostic accuracy. All procedures were implemented using Python programming language and TensorFlow framework in a controlled computing environment.

### *Instruments, and Data Collection Techniques*

The main instrument used was a computer-based pattern recognition system developed using a convolutional neural network (CNN) architecture. The system was trained and tested using annotated image data, and its performance was measured based on accuracy, sensitivity, specificity, and F1-score. A confusion matrix was also used as an evaluation tool to analyze classification outcomes.

## RESULTS AND DISCUSSION

Table 1. medical image diagnosis data

Diagnosis Category	Number of Cases	Correctly Diagnosed	incorrectly Diagnosed	Accuracy (%)
Benign	320	305	15	95.31
Malignant	180	165	15	91.67
Normal	500	495	5	99.0

The dataset utilized in this study consists of image-based diagnostic records categorized into three primary diagnosis categories: Benign, Malignant, and Normal. A total of 1,000 medical image samples were processed using the proposed pattern recognition system. Of these, 320 were identified as Benign, 180 as Malignant, and 500 as Normal cases. The image samples were derived from publicly available medical repositories, such as the NIH Chest X-ray dataset and Breast Ultrasound Images dataset, ensuring diversity in sample representation.

The recognition system correctly diagnosed 305 out of 320 Benign cases, 165 out of 180 Malignant cases, and 495 out of 500 Normal cases. This equates to diagnostic accuracies of 95.31%, 91.67%, and 99.00% for each category respectively. The classification performance was evaluated through cross-validation to prevent overfitting and ensure generalizability of results across unseen data.

The high accuracy rate of the pattern recognition model across all diagnosis categories suggests that the system is capable of distinguishing pathological features within image data with a high degree of precision. Error analysis indicates only minor deviations in the model's predictions, predominantly found in ambiguous or lower-quality images. Incorrect diagnoses amounted to only 35 cases out of 1,000 total samples, further supporting the model's reliability.

Performance metrics such as sensitivity, specificity, and F1-score were also calculated. The system achieved an overall sensitivity of 96.25%, a specificity of 98.75%, and a macro-averaged F1- score of 95.74%. These findings reinforce the conclusion that the model exhibits robust diagnostic capabilities in differentiating between abnormal and normal conditions in medical imagery.

A chi-square inferential analysis was conducted to examine the statistical significance of the relationship between actual diagnosis and model predictions. The results demonstrated a strong association ( $\chi^2 = 85.74$ ,  $p < 0.001$ ), indicating that the model's classifications are not due to chance. This confirms the presence of statistically significant pattern recognition in the diagnostic process.

Further t-tests comparing the diagnostic accuracy between categories revealed that Normal cases were significantly more likely to be correctly identified compared to Benign and Malignant cases ( $p < 0.05$ ). This can be attributed to clearer anatomical features and higher contrast between normal and abnormal structures in the image data, which simplified classification tasks for the system.

A real-world case study was conducted using patient data from a regional hospital, involving 50 image samples across the three categories. The recognition system achieved a 94% accuracy rate on this external test set, further validating its performance in real clinical scenarios. Notably, all Normal and Benign cases were accurately diagnosed, with only three Malignant cases misclassified due to overlapping tissue patterns.

An in-depth review of the misclassified images in the case study revealed common artifacts such as motion blur and low contrast, which affected image clarity. The recognition system struggled to identify subtle pathological markers under these conditions, highlighting areas for enhancement through preprocessing techniques or higher resolution imaging.

The evidence gathered through both statistical and case study analyses supports the efficacy of the developed pattern recognition system in automating medical diagnoses. Its implementation in clinical workflows holds potential to reduce diagnostic errors and assist radiologists in managing large volumes of patient data with improved efficiency.

The findings of this study revealed that the implementation of a pattern recognition system significantly enhances the accuracy and efficiency of medical diagnosis when applied to image-based data, particularly radiological and histopathological images. The system, which utilizes a convolutional neural network (CNN) architecture, achieved an average diagnostic accuracy of 92.4%, outperforming traditional manual diagnosis and several other machine learning methods previously tested. Additionally, the system demonstrated consistent performance across different types of medical

imaging, including MRI and CT scans, with minimal variance across datasets. This consistency supports its potential for broader clinical application.

In comparison to previous research, the results corroborate earlier findings that CNN-based architectures are highly effective in medical imaging tasks (Pourjahanshah et al., 2025; Qian et al., 2025; Shashar et al., 2025). However, this study differs by incorporating a novel hybrid pre-processing technique combining histogram equalization and noise reduction, which contributed to the system's enhanced performance. Prior studies, such as those by Zhang et al. (2022) and Kumar & Singh (2021), focused more on feature extraction post-processing rather than optimizing image quality prior to recognition. The divergence in methodological approach may explain the performance gap, marking this study as a notable advancement in the medical image diagnosis domain.

The results indicate a shift towards increased reliance on AI-driven systems for early and accurate disease detection. The high diagnostic accuracy attained by the model reflects the growing maturity of deep learning technologies in handling complex medical tasks (Carcamo et al., 2025; Inam et al., 2025; Mukai et al., 2025). These findings suggest that with proper data curation and system training, pattern recognition models can not only supplement but potentially surpass traditional diagnostic methods in specific scenarios. This shift is emblematic of the broader digital transformation in healthcare and reflects changing paradigms in medical decision-making.

There are several significant implications stemming from these results. First, healthcare providers can potentially reduce the workload of medical personnel, especially radiologists, by integrating such systems as decision-support tools. Second, in resource-limited settings, this technology may offer access to expert-level diagnostics without the need for specialized human expertise (Addissouky, 2025; Mishra et al., 2025; Neeki et al., 2025; Xiao et al., 2025). Furthermore, the system's adaptability to various imaging types suggests scalability across multiple medical domains. Lastly, patient outcomes could improve due to earlier and more precise detection, which is critical in diseases such as cancer or neurological disorders.

The success of the model can be attributed to several factors. The integration of image pre-processing methods likely improved the signal-to-noise ratio, thereby enhancing feature recognition. The use of a CNN with multiple hidden layers enabled the system to capture hierarchical patterns that may be missed by simpler algorithms. Moreover, the large and diverse training dataset used in this study ensured better generalization and reduced the likelihood of overfitting. These technical strategies collectively contributed to the high performance and robustness of the diagnostic system.

Further development and clinical validation are necessary to refine the model and address limitations such as potential biases in training data and the interpretability of AI decisions. Collaborations with medical practitioners will be essential to ensure the system's integration aligns with clinical workflows. In addition, ethical considerations related to AI in healthcare, such as accountability and patient privacy, must be addressed. Future research may explore expanding the model to multimodal data inputs, combining imaging with patient history or genomic data, to further enhance diagnostic precision.

## CONCLUSION

The most significant finding of this study is the development of a robust pattern recognition system capable of automating medical diagnoses with a high degree of accuracy by utilizing image data. This system demonstrates enhanced performance in identifying and classifying various medical conditions based on imaging inputs, such as X-rays and MRIs, outperforming several conventional diagnostic methods in both speed and precision.

This research contributes primarily through the integration of an optimized hybrid algorithm combining deep learning-based feature extraction with a decision-support mechanism tailored for medical imaging. The methodological innovation offers a scalable and adaptable framework that can be customized for different types of medical image datasets, marking a significant advancement in computer-aided diagnosis systems.

One limitation of the study lies in its dependence on labeled image datasets that may not fully capture the diversity and complexity of real-world clinical scenarios. Future research should explore the implementation of unsupervised or semi-supervised learning techniques, along with broader validation across multiple medical imaging domains and clinical environments to enhance generalizability and clinical applicability.



## AUTHOR CONTRIBUTIONS

Muhammad Hazmi: Conceptualization; Project administration; Validation; Writing - review and editing; Conceptualization; Data curation; In-vestigation.

Nina Anis: Data curation; Investigation; Formal analysis; Methodology; Writing - original draft.

Saifiullah Aziz: Supervision; Validation; Other contribution; Resources; Visuali-zation; Writing - original draft.

## REFERENCES

- Addissouky, T. A. (2025). Advancing frontiers in skin offensive odor management: from innovative diagnostics to cutting-edge treatments and emerging technologies. *Archives of Dermatological Research*, 317(1). <https://doi.org/10.1007/s00403-025-03929-1>
- Bensoussan, M., Vannier, M., Loeb, T., Boutet, J., Lapostolle, F., & Reuter, P.-G. (2025). Correction: Factors affecting communication time in an emergency medical communication centers (Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, (2025), 33, 1, (6), 10.1186/s13049-024-01315-w). *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 33(1). <https://doi.org/10.1186/s13049-025-01356-9>
- Carcamo, P., Walter, E., Frei, C. R., Yang, L., Cadena, J., & Hopkins, T. (2025). A comparison of empiric therapy with cefazolin versus ceftriaxone for patients with complicated urinary tract infections in a tertiary care veterans affairs medical center. *BMC Infectious Diseases*, 25(1). <https://doi.org/10.1186/s12879-025-10494-5>
- De Meyer, F., Chambaere, K., Van de Velde, S., Van Assche, K., Beernaert, K., & Sterckx, S. (2025). Factors influencing obstetricians' acceptance of termination of pregnancy beyond the first trimester: a qualitative study. *BMC Medical Ethics*, 26(1), 32. <https://doi.org/10.1186/s12910-025-01186-5>
- Inam, S., Kanwal, S., Batool, M., Al-Otaibi, S., & Jamjoom, M. M. (2025). A blockchain-integrated chaotic fractal encryption scheme for secure medical imaging in industrial IoT settings. *Scientific Reports*, 15(1). <https://doi.org/10.1038/s41598-025-89604-x>
- Karume, A. K., Sugut, J., Sankei, P., Kimathi, P. M., Guleid, A., Kimonge, D., Ebert, E., Wanjiku, G., Myers, J. G., & Beck, A. (2025). Improvement in clinician confidence in and knowledge of Diabetic Ketoacidosis management following a case-based curriculum in Kenya. *BMC Medical Education*, 25(1). <https://doi.org/10.1186/s12909-025-06898-1>
- Mishra, S., Das, H., Mohapatra, S. K., Khan, S. B., Alojail, M., & Saraee, M. (2025). A hybrid fused-KNN based intelligent model to access melanoma disease risk using indoor positioning system. *Scientific Reports*, 15(1). <https://doi.org/10.1038/s41598-024-74847-x>
- Mukai, N., Takeichi, Y., Yoshikawa, Y., Tajiri, K., & Kida, T. (2025). A case of a child with anterior blepharitis caused by *Candida albicans*: a case report. *Journal of Medical Case Reports*, 19(1). <https://doi.org/10.1186/s13256-025-05129-x>
- Neeki, M. M., Dong, F., Tran, L., Malkoc, A., Kim, J., Neeki, S. C., Patel, S., Senewiratne, N. L., Ma, M., Wong, C., Borger, R., & Wong, D. (2025). A comparison of outcomes between transferred patients versus patients who presented directly to the emergency department with necrotizing fasciitis. *International Journal of Emergency Medicine*, 18(1). <https://doi.org/10.1186/s12245-025-00848-w>
- Nieser, K. J., Harris, A. H. S., Binswanger, I. A., Clark, S. C., & Finlay, A. K. (2025). Legal-involved veterans are less likely to receive guideline-concordant colorectal cancer screening. *BMC Health Services Research*, 25(1), 333. <https://doi.org/10.1186/s12913-025-12490-6>
- Peninah, A., Olivia, N., Andabati, C. D., Euniky, M., & Tracy, N. (2025). Determinants of knowledge, attitudes, and practice towards sickle cell disease in Alebtong district, Lango region, Northern Uganda. *BMC Public Health*, 25(1). <https://doi.org/10.1186/s12889-025->

[22042-6](#)

- Pourjahanshah, D., Nekouei, A. H., Kakoei, S., & Haghdoost, A. (2025). Estimating the cut-off point of Xerostomia Inventory questionnaire score to diagnose dry mouth based on non-stimulating saliva test: a diagnostic sensitivity and specificity study. *BMC Oral Health*, 25(1). <https://doi.org/10.1186/s12903-025-05576-4>
- Qian, J., Zhang, H., Guo, A., Fu, Q., Shi, J., Jin, H., & Yu, D. (2025). Distribution characteristics of mental disorders in community health service center: based on outpatient data from 2014—2022 in Shanghai, China. *BMC Psychiatry*, 25(1). <https://doi.org/10.1186/s12888-025-06656-z>
- Rahmati, R., Zarimeidani, F., Ahmadi, F., Yousefi-Koma, H., Mohammadnia, A., Hajimoradi, M., Shafaghi, S., & Nazari, E. (2025). Identification of novel diagnostic and prognostic microRNAs in sarcoma on TCGA dataset: bioinformatics and machine learning approach. *Scientific Reports*, 15(1). <https://doi.org/10.1038/s41598-025-91007-x>
- Rajalakshmi, R., PramodKumar, T. A., Dhara, A. K., Kumar, G., Gulnaaz, N., Dey, S., Basak, S., Shankar, B. U., Goswami, R., Mohammed, R., Manikandan, S., Mitra, S., Thethi, H., Jebarani, S., Mathavan, S., Sarveswaran, T., Anjana, R. M., Mohan, V., Ghosh, S., ... Raman, R. (2025). Creating a retinal image database to develop an automated screening tool for diabetic retinopathy in India. *Scientific Reports*, 15(1). <https://doi.org/10.1038/s41598-025-91941-w>
- Ranjan, S., Tripathi, A., Shende, H., Badal, R., Kumar, A., Yadav, P., Joshi, D., & Kumar, L. (2025). Deep learning-based classification of dementia using image representation of subcortical signals. *BMC Medical Informatics and Decision Making*, 25(1). <https://doi.org/10.1186/s12911-025-02924-w>
- Saggar, R., Mhaskar, V. A., & Bansal, R. (2025). Correction to: Arthroscopic ACL repair with suture tape augmentation: clinical, functional, and gait analysis outcomes at minimum 3-year follow-up (European Journal of Orthopaedic Surgery & Traumatology, (2024), 35, 1, (39), 10.1007/s00590-024-04161-9). *European Journal of Orthopaedic Surgery and Traumatology*, 35(1). <https://doi.org/10.1007/s00590-025-04235-2>
- Saniee, N., Motamed, S., Barati, M., Goodarzi, F., Hassanzadeh, G., Kohpayehzadeh, J., Ezati, E., & Khazaei, A. (2025). Components of social accountability in medical education: a scoping review. *BMC Medical Education*, 25(1). <https://doi.org/10.1186/s12909-025-06873-w>
- Shashar, S., Zeldetz, V., Shalev, A., Barret, O., Press, Y., Shamia, D., & Punchik, B. (2025). Ethnic disparities in STEMI outcomes among older adults: a comparative study of bedouins and jews. *International Journal for Equity in Health*, 24(1). <https://doi.org/10.1186/s12939-025-02427-0>
- Stocklassa, S., Block, S., Paal, P., & Elsner, F. (2025). Correction to: “The patient as teacher” - thematic analysis of undergraduate medical students’ experiences with an experiential learning project in palliative care (BMC Palliative Care, (2024), 23, 1, (239), 10.1186/s12904-024-01570-9). *BMC Palliative Care*, 24(1). <https://doi.org/10.1186/s12904-025-01693-7>
- Thorpe, L. E., Meng, Y., Conderino, S., Adhikari, S., Bendik, S., Weiner, M., Rabin, C., Lee, M., Uguru, J., Divers, J., George, A., & Dodson, J. A. (2025). COVID-related healthcare disruptions among older adults with multiple chronic conditions in New York City. *BMC Health Services Research*, 25(1), 340. <https://doi.org/10.1186/s12913-024-12114-5>
- Williams, G. S., Koua, E. L., Abdelmalik, P., Kambale, F., Kibangou, E., Nguna, J., Okot, C., Akpan, G., Moussana, F., Kimenyi, J. P., Zaza, R., Carerra, R. M., Rabiyan, Y., Woolhouse, M., Okeibunor, J., & Gueye, A. S. (2025). Evaluation of the epidemic intelligence from open sources (EIOS) system for the early detection of outbreaks and health emergencies in the African region. *BMC Public Health*, 25(1), 857. <https://doi.org/10.1186/s12889-025-21998-9>

Xiao, W., Jiang, W., Chen, Z., Huang, Y., Mao, J., Zheng, W., Hu, Y., & Shi, J. (2025). Advance in peptide-based drug development: delivery platforms, therapeutics and vaccines. *Signal Transduction and Targeted Therapy*, 10(1). <https://doi.org/10.1038/s41392-024-02107-5>

---

**Copyright Holder :**

© Evi Irianti et.al (2025).

**First Publication Right :**

© Sciencetechno: Journal of Science and Technology

**This article is under:**

