

The Relationship Between Knowledge About Kangaroo Mother Care (Kmc) Among Mothers and Increased Weight of Low Birth Weight (Lbw) Babies in the Neonate Intensive Care Room of the Mayapada Hospital Jakarta Selatan

Nuke Tri Indriani¹, Arozamati W. Lase², Rome Tao Toba Muara Ria³, Sicily Prima Yanuaria Buka⁴, Herlina Meilina⁵

¹ Mayapada Hospital Jakarta Selatan, Indonesia

² Universitas Borobudur, Indonesia

³ Universitas Borobudur, Indonesia

⁴ Mayapada Hospital Jakarta Selatan, Indonesia

⁵ Mayapada Hospital Jakarta Selatan, Indonesia

Corresponding Author:

Nuke Tri Indriani,

Mayapada Hospital Jakarta Selatan, Indonesia

Jl. Lebak Bulus I No. Kav. 29, RT.6/RW.4, Lb. Bulus, Kec. Cilandak, Kota Jakarta Selatan, Daerah Khusus Ibukota Jakarta 12440

Email: nukeindriani79@gmail.com

Article Info

Received: Sep 10, 2024

Revised: Nov 5, 2024


Accepted: Dec 2, 2024

Online Version: Feb 6, 2025

Abstract

One of the main causes of infant mortality is low birth weight babies. Low birth weight babies are high risk babies and are susceptible to various problems such as the risk of respiratory infections, low blood sugar levels (hypoglycemia), eating disorders, too many red blood cells and the risk of experiencing hypothermia. In general, babies with low birth weight must be treated with an incubator or using the KMC method so that their body temperature is maintained. This study aims to determine the relationship between knowledge of the Kangaroo Mother Care (KMC) method among mothers and the increase in Low Birth Weight (LBW) Babies in the NICU Room at Mayapada Hospital, South Jakarta. This research is quantitative using a correlative analytical descriptive design with a cross sectional approach. Shows that there is a significant relationship between knowledge about the Kangaroo Mother Care (KMC) Method in mothers and the increase in Low Birth Weight (LBW) in the NICU room at Mayapada Hospital, South Jakarta.

Keywords: Kangaroo, Knowledge, LBW



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/health> ISSN: (P: 2988-7550) - (E: 2988-0459)

How to cite:

Indriani, T, N., Lase, W, A., Ria, M, T, T, R., Buka, Y, P, S & Meilina, H. (2025). The Relationship Between Knowledge About Kangaroo Mother Care (Kmc) Among Mothers and Increased Weight of Low Birth Weight (Lbw) Babies in the Neonate Intensive Care Room of the Mayapada Hospital Jakarta Selatan. *Journal of World Future Medicine, Health and Nursing*, 3(1), 18–28. <https://doi.org/10.70177/health.v3i1.1862>

Published by:

Yayasan Adra Karima Hubbi

INTRODUCTION

Data from the World Health Organization (WHO) in 2018 states that the prevalence of LBW babies in the world is 15.5% or around 20 million are born every year, around 96.5% of which occur in developing countries. One of the highest contributors to the Neonatal Mortality Rate (AKN) is LBW (Kazumova & Samburova, 2025; Mursil et al., 2025). The incidence of LBW in the world is currently 15.5% (which means around 20.6 million every year), and 96.5% in developing countries. Indonesia is one of the countries in Southeast Asia with a LBW prevalence of more than 15.5% of births every year. In 2018 the results of Riskesdas showed that the proportion of body weight less than 2500 grams in Indonesia on average was 6.2% (Juza et al., 2025; Wiegel et al., 2025).

Low birth weight babies are high risk babies and are susceptible to various problems such as the risk of respiratory infections, low blood sugar levels (hypoglycemia), eating disorders, too many red blood cells and are at risk of experiencing hypothermia due to difficulties in adapting to the environment outside the womb after birth (Balcells-Esponera et al., 2025; Wang et al., 2025). This is due to the immaturity of the baby's organ systems such as the lungs, kidneys, heart, immune system and digestive system. Babies with low birth weight will experience hypothermia caused by very thin subcutaneous fat so it is influenced by environmental temperature. In general, babies with low birth weight must be treated with an incubator or use the KMC method so that their body temperature is maintained (May et al., 2025; Okonogi et al., 2025). Several perinatologists conducted several studies and concluded that care using the kangaroo method or what is called the attachment method can provide more benefits in treating babies weighing less than 2,500 grams (Hammond et al., 2025; Oksuzoglu & Okumus, 2025).

The results of a preliminary study that has been carried out, from medical record data from Mayapada Hospital, South Jakarta from June to August 2023, showed data on LBW babies in the NICU room as many as 54 babies with details of hyperthermia as many as 8 babies (14.8%), hypothermia as many as 20 babies (37. %) and normal temperature for 26 babies (48.2%). The results of observations carried out by researchers, from the birth data of 54 LBW babies, only 15 babies (27.8%) were carried out by KMC. Ten mothers (20.4%) performed baby care using the KMC method for less than one hour and eight mothers (14.8%) performed KMC on their babies for approximately one hour. Based on the background above, researchers were encouraged to conduct research with the title Relationship between knowledge of the Kangaroo Mother Care (KMC) method and an increase in Low Birth Weight (LBW) in the NICU Room at Mayapada Hospital, South Jakarta.

RESEARCH METHOD

This study was conducted to examine the relationship between maternal knowledge of Kangaroo Mother Care (KMC) and infant weight gain using a quantitative approach. The research was carried out over a period from October 2023 to the end of February 2024. Ethical considerations were strictly maintained throughout the study, including respect for human dignity through informed consent, anonymity of respondents, confidentiality of data, and adherence to principles of fairness and transparency. These ethical standards ensured that participants were protected and that the research process remained credible and accountable.

Research Design

This research employed a descriptive analytical design with a cross-sectional approach, aiming to analyze variables at a single point in time. The study sought to describe and examine the association between mothers' knowledge of KMC and changes in infant weight. Sampling was conducted using a total sampling technique, where all eligible participants who met the inclusion and exclusion criteria were included in the study. This design allowed for a systematic and structured analysis of the relationship between the variables under investigation (Goll et al., 2025; Saberian et al., 2025).

Research Target/Subject

The subjects of this study consisted of mothers who met predetermined inclusion and exclusion criteria. By applying the total sampling method, all accessible and qualified respondents were included as research participants. This approach ensured comprehensive data representation and minimized sampling bias, allowing the findings to more accurately reflect the characteristics of the target population involved in KMC practices.

Research Procedure

The research procedure began with obtaining ethical approval and informed consent from participants. Prior to data collection, the research instruments underwent validity and reliability testing using IBM SPSS version 27 software. After confirming the instruments were appropriate, respondents were asked to read and complete the questionnaire. In addition, infant weight measurements were conducted twice, during the first and second weeks, using standardized procedures to observe any changes in weight. These steps were carried out systematically to ensure consistency and accuracy of the data collected.

Instruments and Data Collection Techniques

Data collection in this study utilized several instruments, including a knowledge questionnaire consisting of 20 true-or-false questions, documentation sheets for recording infant weight, and a baby scale measured in grams. The questionnaire had previously undergone validity testing, yielding correlation values ranging from 0.581 to 0.919, exceeding the r-table value of 0.361, indicating that all items were valid. Reliability testing showed a Cronbach's Alpha value of 0.786, categorizing the instrument as highly reliable. Data were collected through questionnaire distribution, documentation studies, and direct observation. Infant weight was measured with the baby in a supine position, ensuring accuracy during the measurement process.

Data Analysis Technique

Data analysis was conducted using quantitative techniques, including data editing, coding, scoring, and tabulation. The analysis involved both univariate and bivariate approaches. Univariate analysis was used to generate frequency distributions of the studied variables, while bivariate analysis employed the Chi-Square test to examine the relationship between maternal knowledge of KMC and infant weight gain. The level of significance was determined using a p-value threshold of 0.05. If the p-value was less than or equal to 0.05, the null hypothesis was rejected, indicating a significant relationship between variables. Conversely, if the p-value exceeded 0.05, the null hypothesis was accepted, indicating no significant relationship between the variables studied.

RESULTS AND DISCUSSION

The research results related to the characteristics of the respondents in this study, namely:

Characteristics of mothers of LBW babies in the MHJS nursery

Age frequency distribution

Univariate analysis of this study includes the characteristics of respondents, age, parity, and pregnancy history.

Age Frequency Distribution in the NICU Room at Mayapada Hospital, South




Figure 1. Age frequency distribution in the nicu room at mayapada hospital, south jakarta

The research results showed that the majority of respondents who had babies were middle adults (21-35 years) amounting to 56.7% or 17 respondents (n= 30). In this study, it was found that the age was in accordance with the applicable theory. Age 20-35 years is the ideal age for a mother to become pregnant and give birth because at that age the mother's uterus is mature and able to accept pregnancy well from a mental and physical perspective. According to Qurniyawaty et al (2014), maternal age 20-35 years during pregnancy has the lowest risk of pregnancy and childbirth. This means that a mother over 35 years of age at the time of pregnancy has a very high risk of pregnancy and childbirth which is detrimental to the health of the mother and the child to be born.

In addition, maternal death at the age of <20 years and >35 years is an age at risk that is 2-5 times higher than at the age of 20-35 years. Apart from that, according to Pinontoan (2015), the relationship between maternal age and the incidence of LBW is due to the fact that at <20 years of age the development of the reproductive system is not yet optimal so it affects the birth weight of the baby, whereas in mothers aged >35 years the function of the reproductive system has decreased so that it will affect the pregnancy. Also, as the mother ages, there will be changes in the blood vessels and a decrease in the function of hormones that regulate the reproductive cycle. If the mother's age is within the age range that is not at risk then the chance of LBW will also be low, conversely if the mother is of high risk age then the chance of LBW will be higher.

Parity frequency distribution




Figure 2. Parity frequency distribution in the NICU room at mayapada hospital, south jakarta

Figure 2 Distribution shows that the majority of respondents' parity is between 2 and 3 children, 56.7% or 17 respondents. High parity will have an impact on various health outcomes for both the mother and the baby born. Repeated pregnancies and childbirth will cause damage to the blood vessels in the walls of the uterus and a decline in the flexibility (elasticity) of the tissue that has been repeatedly stretched during pregnancy so that there tends to be abnormalities in the location or growth of the placenta and fetal growth, resulting in the birth of babies with low birth weight (Pinontoan, 2015).

Frequency Distribution of Pregnancy History




Figure 3. Frequency distribution of pregnancy history in the NICU room at mayapada hospital, south jakarta

Diagram 3 above shows that all respondents' pregnancy history is not 100% multiple. Biologically, multiple pregnancies are at risk of LBW. In multiple pregnancies, the weight of one multiple fetus is on average 1000 grams lighter than a single fetus. The weight of the fetuses from multiple pregnancies is not the same, generally, there is a difference of between 50 and 1000 grams. Apart from that, there is an unequal distribution of blood circulation. As a result, the growth of the two fetuses was different. Multiple pregnancies can pose a higher risk

to the baby and mother. The need for growth in multiple pregnancies is greater so that if nutritional deficiencies such as pregnancy anemia occur, it can disrupt fetal growth in the womb (Sulistiani, 2014).

Frequency distribution of KMC knowledge among mothers




Figure 4. Frequency distribution of kmc knowledge among mothers in the nicu room at mayapada hospital, south jakarta

Figure 4 shows that almost all respondents' knowledge about KMC is in the good category, amounting to 76.7% or 23 respondents. According to Siswandoyo (2016), knowledge can be defined as a collection of information that can be understood and obtained from the learning process throughout life and can be used at any time as a tool for self-adjustment. Knowledge is an introduction to reality, truth, principles and rules of an object and is the result of information stimulation for behavior change. The knowledge that mothers need to understand regarding the kangaroo mother care method includes mothers needing to know the meaning of KMC, the benefits of KMC for LBW babies, how to carry out KMC and the criteria for the success of KMC. Once the mother has good knowledge about KMC, the mother can then determine the attitudes and behavior that can be taken for the welfare of her baby (Sofiana J, et al. 2016). The results of this research are in line with research from Kurniawati et al., (2022).

The research results showed that respondents had good knowledge (46.7%), sufficient knowledge (20.0%), and poor knowledge (33.3%). Other research results from (Amalia & Herawati, 2018) showed that some respondents had good knowledge, namely 34 mothers (45.3%) and those with sufficient knowledge criteria were 14 mothers (18.7%) and those with insufficient knowledge criteria were 27 mothers (36%). Research results that are not in line with Suwarni et al., (2021) as many as 7 respondents (35%) had good knowledge and 13 respondents (65%) had poor knowledge. In fact, taking KMC for a longer period of time has a positive effect on the length of breastfeeding and the baby's temperature within the normal range, resulting in an increase in the baby's weight. A baby who breastfeeds the mother for longer will make the baby feel calm and comfortable so that the baby gets an adequate supply of breast milk and the energy obtained by the body is only focused on growth.

Frequency Distribution of Increased Weight in LBW babies

Shows that almost all respondents' knowledge about KMC is in the good category, amounting to 76.7% or 23 respondents. Babies with low birth weight (LBW) are those whose body weight is less than 2,500 grams regardless of the gestation period. LBW is a major factor in increasing mortality, morbidity and disability of neonates, infants and children and has a long-term impact on their lives in the future. Therefore, it is necessary to provide care for babies with LBW (Sembiring, 2017).

The increase in body weight also occurs due to the increased relationship between the baby and the mother, where the baby has more time to feel touch, which can reduce the release of catecholamines in the blood, thereby reducing the physiological stress of the fetus. KMC increases mother and baby bonding and is a therapeutic intervention to increase maternal closeness, promote natural behavior to stimulate growth and development (Arifah & Wahyuni, 2016).

The results of this research are (Amalia & Herawati, 2018) that the implementation of Kangaroo Method Care (PMK) is influenced by the knowledge and attitudes of mothers who have Low Birth Weight babies with a p value of $0.004 < (0.05)$. Other research from Gratisari & Agonwardi (2017) was conducted at RSUD Dr. Rasidin on 15 mothers who have LBW. The results of the study showed a p value of 0.001, which means that there is an influence of KMC on weight gain in LBW. Apart from that, research from (Munns et al., 2025; Rocha et al., 2025) also concluded that KMC is effective in increasing the weight of LBW babies.

The relationship between knowledge of the kangaroo mother care (KMC) method in mothers and increased weight in babies with low birth weight (LBW)

Chi Square Test Results of the Relationship between Knowledge of the Kangaroo Mother Care (KMC) Method and Weight Gain in Low Birth Weight (LBW) Babies in the Mayapada Hospital Nicu Room, South Jakarta

Table 1. Chi square test results of the relationship

Knowledge	Weight Gain	OR	P-Value
	Increase n	Still %	Total n
Good	23	76.8	0

The results of the crosstabulation test between knowledge and weight gain in LBW babies showed that all respondents with good knowledge experienced an increase in body weight in LBW babies (100%). Meanwhile, most respondents with less knowledge did not experience weight gain in LBW babies (57.1%). The Chi Square test results show that the p value is $0.001 < 0.05$ with an OR value of 6.5, which means that H_a is accepted and H_0 is rejected. It can be concluded that there is a relationship between knowledge of the Kangaroo Mother Care (KMC) Method and weight gain in babies with low birth weight (LBW) in the NICU room at Mayapada Hospital, South Jakarta, and mothers with good knowledge have a 6 times greater chance of increasing weight. body weight in their babies, while mothers with less knowledge have a chance of having babies with a constant weight.

According to Notoatmodjo, (2018) that knowledge is a domain that is quite important in determining behavior. Behavior that is based on knowledge, awareness and positive attitudes will be more lasting. Good knowledge will make it easier for someone to change behavior, including KMC practices (Ananda & Nangia, 2025; Saberian et al., 2025). KMC can increase

the weight and body length of LBW babies. KMC is highly recommended for LBW babies because when the mother's skin meets the baby's skin there will be a change in temperature. This happens because the baby is relaxed, resting in a pleasant position, similar to the position in the womb, so that the baby is less anxious and sleeps longer. In this situation, oxygen and calorie consumption is at the lowest level, so the existing calories are used to increase body weight. In addition, the baby's weight gain is caused by increased breast milk production and more frequent breastfeeding, which is one of the benefits of KMC.

The results of this research are in line with research from Kurniawati et al., (2022) The research results showed that respondents had good knowledge (46.7%), sufficient knowledge (20.0%), and poor knowledge (33.3%). The results of the analysis of the relationship between maternal knowledge and the implementation of KMC showed p -value = 0.012, which means there is a relationship between maternal knowledge and the implementation of the Kangaroo Mother Care (KMC) method for low birth weight (LBW) babies. Other research in line from Wulaningsih et al., (2023) that the research results show that there is an influence of KMC on increasing LBW body weight through a maximum increase in the graph from day to day 5 in 2012, amounting to 88 grams. The test results obtained a p value (0.000) and comparing the weight when weighed with the weight at start showed a significant average of -93.576 (CI - 220.376/-126.8) and a standard deviation of 91.00.

CONCLUSION

Based on the research results, it shows that: The majority of respondents who have babies are 21-35 years old, 56.7%, the parity of respondents with 2 and 3 children is 56.7%, and all of the respondents' pregnancy history is not multiple (100%). Most of the respondents' knowledge about KMC was in the good category at 76.7%. Most of the weight gain in babies with LBW increased by 86.7%. There is a relationship between knowledge of the Kangaroo Mother Care (KMC) Method in mothers and weight gain in babies with Low Birth Weight (LBW) in the NICU Room at Mayapada Hospital, South Jakarta with a p value of $0.001 < 0.05$.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration.

Author 2: Data curation; In-vestigation.

Author 3: Validation; Writing.

Author 4: Review and Editing.

Author 4: Formal analysis; Methodology; Writing - original draft.

Author 5: Conceptualization; Data curation; In-vestigation.

REFERENCES

- Ananda, R., & Nangia, S. (2025). Influence of Early Total Enteral Feeding in Preterm Infants with Respiratory Distress Syndrome. *Neonatology*, 122(1), 4–10. Scopus. <https://doi.org/10.1159/000539544>
- Balcells-Esponera, C., Aldecoa-Bilbao, V., Borrás Novell, C., López-Abad, M., Valls-Lafon, A., Batllori Tragant, M., Izquierdo-Renau, M., del Rey Hurtado de Mendoza, B., Herranz-Barbero, A., & Iglesias-Platas, I. (2025). Intestinal Biomarkers in Preterm Infants: Influence of Mother's Own Milk on Fecal Calprotectin and of Gestational Age on IFABP Concentrations. *Nutrients*, 17(13). Scopus. <https://doi.org/10.3390/nu17132177>

- Belachew, E. A. (2023). Perception, willingness, and practices of telemedicine in patients with chronic diseases: Implication of digital health in patients' perspective at a tertiary care hospital in Ethiopia. *Frontiers in Public Health*, *11*(Query date: 2025-02-02 19:25:32). <https://doi.org/10.3389/fpubh.2023.1234436>
- Bilal, W. (2022a). Digital health and telemedicine in Pakistan: Improving maternal healthcare. *Annals of Medicine and Surgery*, *81*(Query date: 2025-02-02 19:25:32). <https://doi.org/10.1016/j.amsu.2022.104425>
- Bilal, W. (2022b). Improving access to maternal care in Africa through telemedicine and digital health. *International Journal of Health Planning and Management*, *37*(4), 2494–2500. <https://doi.org/10.1002/hpm.3498>
- Boikanyo, K. (2023). Remote patient monitoring systems: Applications, architecture, and challenges. *Scientific African*, *20*(Query date: 2025-02-02 12:26:20). <https://doi.org/10.1016/j.sciaf.2023.e01638>
- Bolam, S. M. (2021). Remote patient monitoring with wearable sensors following knee arthroplasty. *Sensors*, *21*(15). <https://doi.org/10.3390/s21155143>
- Borrelli, N. (2023). Telemedicine in Adult Congenital Heart Disease: Usefulness of Digital Health Technology in the Assistance of Critical Patients. *International Journal of Environmental Research and Public Health*, *20*(10). <https://doi.org/10.3390/ijerph20105775>
- Bruining, N. (2021). The post-pandemic legacy: The breakthrough of digital health and telemedicine. *Cardiovascular Research*, *117*(9). <https://doi.org/10.1093/cvr/cvab178>
- de Lima Castro, M., Dos Passos, R. R., Rocha, T. L., do Amaral, W. N., & Giachini, F. R. (2025). Increased protein O-GlcNAcylation in human placentas from hypertensive pregnancies. *Placenta*. Scopus. <https://doi.org/10.1016/j.placenta.2025.05.024>
- Donkor, D. R., Ziblim, S.-D., Asumah, M. N., Dzantor, E. K., & Abdul-Mumin, A. (2025). Incidence and risk factors for neonatal jaundice in a teaching hospital in Northern Ghana. *BMC Pediatrics*, *25*(1). Scopus. <https://doi.org/10.1186/s12887-025-06221-3>
- Goll, N., Wiechers, C., Zinke, K., Preißl, H., Fritsche, A., & Hallschmid, M. (2025). Influence of maternal metabolism on neonate eating behaviour. *Appetite*, *214*. Scopus. <https://doi.org/10.1016/j.appet.2025.108206>
- Hammond, S. J., Roff, A. J., Robinson, J. L., Darby, J. R. T., Meakin, A. S., Clifton, V. L., Bischof, R. J., Stark, M. J., Wallace, M. J., Tai, A., Morrison, J. L., & Gatford, K. L. (2025). In utero exposure to experimental maternal asthma alters fetal airway development in sheep. *Experimental Physiology*, *110*(6), 899–907. Scopus. <https://doi.org/10.1113/EP092502>
- Hillick, D., O'Reilly, D., Murphy, L., Breathnach, F., & McCallion, N. (2025). Increased risk of admission to neonatal intensive care unit in neonates born to mothers with pregestational diabetes. *European Journal of Pediatrics*, *184*(6). Scopus. <https://doi.org/10.1007/s00431-025-06170-0>
- Jana, A., Pramanik, M., Maiti, A., Chattopadhyay, A., & Abed Al Ahad, M. A. (2025). In-utero exposure to PM2.5 and adverse birth outcomes in India: Geostatistical modelling using remote sensing and demographic health survey data 2019–21. *PLOS Global Public Health*, *5*(7 July). Scopus. <https://doi.org/10.1371/journal.pgph.0003798>
- Juza, A., Kołodziej-Spirodek, L., & Dąbrowski, M. (2025). Important Role of Pregnancy Planning in Pregnancy Outcomes in Type 1 Diabetes. *Diabetology*, *6*(8). Scopus. <https://doi.org/10.3390/diabetology6080075>
- Kazumova, A. B., & Samburova, N. V. (2025). Inherited disorders of hemostasis in the pathogenesis of obstetric pathology. *RUDN Journal of Medicine*, *29*(1), 124–134. Scopus. <https://doi.org/10.22363/2313-0245-2025-29-1-124-134>
- May, M. F., McKinney, M. L., Cestare, D., Hussey, A., Zevallos, A. B., Garber, S., Posencheg, M. A., Puopolo, K. M., & Mukhopadhyay, S. (2025). Improving Time to First Feeding

- for Preterm Infants: A Quality Improvement Approach. *Pediatric Quality and Safety*, 10(2). Scopus. <https://doi.org/10.1097/pq9.0000000000000798>
- Marlina, D., Utomo, A., Adriansyah, P. N. A., Pelitawati, D. R., Poernomo, M. A. P. D., Sumawan, H., Handono, B., & Aziz, M. A. (2025). Association of Bacteriuria with Hypertension Risk in Pregnant Women. *Medical Science Monitor*, 31. Scopus. <https://doi.org/10.12659/MSM.946167>
- Marlina, D., Utomo, A., Handono, B., Pelitawati, D. R., Adriansyah, P. N. A., Aziz, M. A., & Aditiyono, A. (2024). Association Between Pre-Pregnancy Body Mass Index and Labor Induction Success Rates: A Case Control Study. *Medical Science Monitor*, 30. Scopus. <https://doi.org/10.12659/MSM.946357>
- Moegis, F., Yudianto, A., & Sumino, R. (2024). Cerebral Vessel Rupture Leading to Sudden Death: Implications for Pharmacognosy. *Pharmacognosy Journal*, 16(6), 1442–1446. Scopus. <https://doi.org/10.5530/pj.2024.16.233>
- Mulyanto, J., Wibowo, Y., Ernawati, D. A., & Lestari, D. W. D. (2024). Assessing inequalities in the hypertension management under national health insurance: Evidence from Southern Central Java, Indonesia. *Journal of Public Health and Development*, 22(2), 180–192. Scopus. <https://doi.org/10.55131/jphd/2024/220215>
- Munns, L. B., Noonan, M., Romano, D. L., & Preston, C. E. J. (2025). Interoceptive and exteroceptive pregnant bodily experiences and postnatal well-being: A network analysis. *British Journal of Health Psychology*, 30(3). Scopus. <https://doi.org/10.1111/bjhp.70002>
- Mursil, M., Rashwan, H. A., Khalid, A., Cavallé-Busquets, P., Santos-Calderón, L., Murphy, M. M., & Puig, D. (2025). Interpretable deep neural networks for advancing early neonatal birth weight prediction using multimodal maternal factors. *Journal of Biomedical Informatics*, 166. Scopus. <https://doi.org/10.1016/j.jbi.2025.104838>
- Nadhiroh, S. R., Fitria, A. L., Hasugian, A. R., Triyono, E. A., Nugroho, N. T., Sinatrya, A. K., & Majid, H. B. A. (2024). Body mass index as the main predictor for length of stay in COVID-19 patients with mild and moderate symptoms: A cross-sectional study in COVID-19 emergency hospital in Indonesia. *Jurnal Ners*, 19(2), 165–172. Scopus. <https://doi.org/10.20473/jn.v19i2.53973>
- Nilansari, A. F., Wahid, R. A. H., & Fatimah, F. A. (2024). Clinical outcomes: Analysis of quality of life and direct medical costs of outpatient hypertensive patients at Panembahan Senopati Hospital, Indonesia. *Journal of Public Health and Development*, 22(1), 145–156. Scopus. <https://doi.org/10.55131/jphd/2024/220111>
- Okonogi, S., Annan, R. A., & Sakurai, T. (2025). Improving Infant Nutrition Through the Market: Experimental Evidence from Ghana. *Journal of Development Studies*, 61(6), 951–969. Scopus. <https://doi.org/10.1080/00220388.2024.2441730>
- Oksuzoglu, M. E., & Okumus, H. G. (2025). Impulsivity, internalizing symptoms, and disordered eating as predictors of adolescents' body mass index: A structural equation modeling study in adolescents with overweight and obesity and their mothers. *Current Psychology*, 44(11), 11170–11185. Scopus. <https://doi.org/10.1007/s12144-025-07956-Z>
- Rocha, A. S., Rangel Bousquet Carrilho, T., Costa, P. R. F., Paixao, E. S., Silva, N. J., da Silva, H. B. M., Falcao, I. R., Fiaccone, R. L., Barreto, M. L., & Ribeiro-Silva, R. D. C. (2025). Interpregnancy Weight Change and Adverse Birth Outcomes: Cohort Study Using Brazil's Routine Register-Based Linked Data. *Maternal and Child Nutrition*, 21(4). Scopus. <https://doi.org/10.1111/mcn.70052>
- Saberian, S., Gale, C., Subhedar, N., Gallagher, N., Esan, O. B., Sinha, I., Harvey, K., Schlüter, D. K., & Taylor-Robinson, D. (2025). Inequalities in neonatal unit mortality in England and Wales between 2012 and 2022: A retrospective cohort study. *The Lancet Child and*

- Adolescent Health*, 9(12), 857–867. Scopus. [https://doi.org/10.1016/S2352-4642\(25\)00243-3](https://doi.org/10.1016/S2352-4642(25)00243-3)
- Talukder, A., Kelly, M., Gray, D., & Sarma, H. (2025). Influence of Maternal Education and Household Wealth on Double Burden of Malnutrition in South and Southeast Asia. *Maternal and Child Nutrition*, 21(4). Scopus. <https://doi.org/10.1111/mcn.70049>
- Wang, X., Kong, X., Ding, Y., An, M., Zhu, X., Guan, Y., & Niu, Y. (2025). Inverted day-night feeding during pregnancy affects the brain health of both maternal and fetal brains through increasing inflammation levels associated with dysbiosis of the gut microbiome in rats. *Journal of Neuroinflammation*, 22(1). Scopus. <https://doi.org/10.1186/s12974-025-03447-x>
- Wiegel, R. E., Baker, K., Calderon-Toledo, C., Gomez, R., Gutiérrez-Cortez, S., Houck, J. A., Larrea, A., Lazo-Vega, L., Moore, L. G., Pisc, J., Toledo-Jaldin, L., & Julian, C. G. (2025). Impaired maternal central hemodynamics precede the onset of vascular disorders of pregnancy at high altitude. *American Journal of Physiology - Heart and Circulatory Physiology*, 328(1), H174–H185. Scopus. <https://doi.org/10.1152/ajpheart.00520.2024>
-

Copyright Holder:

© Nuke Tri Indriani et.al (2025).

First Publication Right:

© Journal of World Future Medicine, Health and Nursing

This article is under:

