

Social Isolation and Cognitive Decline in Aging Populations: AI-Powered Monitoring Systems for Early Detection

Omar Khan¹, Clara Mendes², Marcus Tan³, Ardi Azhar Nampira⁴

¹ Kabul University, Afghanistan

² Universidade Estadual Campinas, Brazil

³ Duke-NUS Medical School, Singapore

⁴ Institut Teknologi Sepuluh November, Indonesia

Corresponding Author:

Omar Khan,
Kabul University, Afghanistan
Kabul, Afghanistan
Email: omarkhan@gmail.com

Article Info

Received: March 2, 2025

Revised: May 7, 2025


Accepted: June 2, 2025

Online Version: Aug 10, 2025

Abstract

Social isolation is a significant and modifiable risk factor for accelerated cognitive decline and dementia in aging populations. Traditional methods for detecting cognitive changes, such as clinical screenings, are often infrequent and fail to capture the subtle, early behavioral shifts that precede a formal diagnosis. This study aimed to develop and validate an artificial intelligence model designed for the early detection of cognitive decline by passively monitoring behavioral and vocal biomarkers of social isolation in older adults living independently. A 24-month, prospective longitudinal study was conducted with a cohort of 200 community-dwelling adults aged 70 and older. A suite of unobtrusive in-home sensors was used to passively collect data on movement patterns, social communication (frequency and duration of conversations), and computer/phone usage. The AI-powered system identified individuals who would later show clinically significant cognitive decline with an accuracy of 91% and a lead time of approximately 7 months before formal assessment. The model successfully distinguished between simple loneliness and the specific behavioral patterns of social withdrawal associated with cognitive impairment. AI-powered passive monitoring systems are a highly effective and ecologically valid tool for the pre-clinical detection of cognitive decline linked to social isolation.

Keywords: Artificial Intelligence, Cognitive Decline, Early Detection



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/health> ISSN: (P: 2988-7550) - (E: 2988-0459)

How to cite:

Khan, O., Mendes, C., Tan, M & Nampira, A, A. (2025). Social Isolation and Cognitive Decline in Aging Populations: AI-Powered Monitoring Systems for Early Detection. *Journal of World Future Medicine, Health and Nursing*, 3(4), 279–292. <https://doi.org/10.70177/health.v3i4.2371>

Published by:

Yayasan Adra Karima Hubbi

INTRODUCTION

The global demographic shift towards an older population presents one of the most significant public health challenges of the 21st century. As longevity increases, so does the prevalence of age-related cognitive decline and neurodegenerative diseases such as Alzheimer's and other dementias. This escalating issue places an immense strain on healthcare systems, families, and economies worldwide. Consequently, the scientific community has intensified its focus on identifying modifiable risk factors that could serve as targets for preventative interventions (Anthonypillailionel & Oruthotaarachchi, 2025; Ba et al., 2025). The goal has shifted from merely treating dementia to proactively preserving cognitive health and promoting successful aging throughout the lifespan.

A compelling and growing body of epidemiological evidence has identified social isolation as a potent and independent risk factor for accelerated cognitive decline. Social isolation, defined as an objective state of having minimal social contact with others, has been robustly linked to poorer cognitive function, a faster rate of cognitive decline, and an increased risk of developing dementia (Pitzalis et al., 2025; Z. Zhang & Zhang, 2025). The mechanisms underlying this association are believed to be multifaceted, involving reduced cognitive stimulation, increased chronic stress and inflammation, and negative impacts on brain structure and function. Unlike genetic predispositions, social isolation is a potentially modifiable factor, making it a prime target for public health interventions aimed at preserving cognitive vitality in older adults.

The convergence of artificial intelligence (AI) and ubiquitous sensing technologies has created an unprecedented opportunity to address this challenge. AI-powered passive monitoring systems, utilizing a network of unobtrusive sensors placed within an individual's home, can continuously and objectively gather data on daily behaviors and routines (Li et al., 2025; Sunith Babu et al., 2025). This technology offers a novel paradigm for health monitoring, moving beyond infrequent, clinic-based assessments to a continuous, real-world understanding of an individual's functional status. In the context of cognitive aging, these systems hold the potential to detect the subtle, early behavioral shifts associated with social withdrawal that may precede clinically apparent cognitive decline.

The primary problem in the early detection of cognitive decline is the profound limitation of current diagnostic methods. Standard clinical practice relies on periodic neuropsychological assessments and patient self-reports (Islam et al., 2025; Li et al., 2025). These assessments are typically conducted only after an individual or their family members have already noticed significant cognitive changes. This approach is inherently reactive rather than proactive, meaning that by the time a diagnosis is made, a substantial degree of irreversible neurodegeneration may have already occurred. This "too little, too late" diagnostic model creates a critical missed window of opportunity for early intervention.

This diagnostic challenge is compounded by the difficulty in accurately and objectively measuring social isolation (Raval et al., 2025; Tyagi, Tiwari, et al., 2025). Traditional methods, such as questionnaires and self-reports, are subject to recall bias, social desirability bias, and a lack of insight from the individual, particularly if cognitive decline has already begun. An older adult may not perceive or wish to admit the extent of their isolation. The specific problem is the absence of a reliable, objective, and continuous method for quantifying the key behavioral biomarkers of social isolation such as changes in communication patterns, time spent outside the home, and engagement in social activities.

The confluence of these issues creates the central problem this research confronts: the lack of a non-invasive, ecologically valid, and scalable system for the pre-clinical detection of cognitive decline. There is a critical need for a tool that can passively and continuously monitor for the subtle behavioral signatures of social withdrawal that are known to be predictive of cognitive impairment (Bataineh et al., 2025; Kogel-Hollacher et al., 2025). Without such a system, the ability to identify at-risk individuals early enough to implement meaningful preventative strategies such as interventions to increase social engagement remains severely limited.

The principal objective of this study is to develop and validate a sophisticated artificial intelligence model capable of the early detection of cognitive decline through the passive, continuous monitoring of in-home behavioral and vocal biomarkers. This research aims to leverage a suite of unobtrusive sensors to collect longitudinal data on daily routines and social interaction patterns (Bataineh et al., 2025; Tyagi, Kumari, et al., 2025). The central goal is to train a recurrent neural network (RNN) to identify subtle changes in these patterns that are predictive of a future decline in cognitive function, as determined by gold-standard neuropsychological assessments.

This study pursues several critical secondary objectives to establish the clinical utility and specificity of the AI model. The first is to identify the most powerful and reliable behavioral predictors of cognitive decline from the multi-modal sensor data. The research seeks to determine which specific biomarkers such as decreased frequency of conversations, increased time spent sedentary, or changes in digital communication are most strongly associated with subsequent cognitive impairment. A second objective is to quantify the “lead time” provided by the AI system, establishing how many months in advance the model can predict a clinically significant cognitive decline compared to traditional assessment methods.

Ultimately, this research aims to produce a robustly validated, AI-powered monitoring system that can serve as a scalable tool for pre-clinical risk stratification in aging populations. The study endeavors to create a system that can distinguish between the behavioral patterns of benign loneliness and the specific, pathological social withdrawal that heralds the onset of cognitive decline (Kumar et al., 2025; Tyagi, Kumari, et al., 2025). The expected outcome is a non-invasive, privacy-preserving technology that can empower individuals, families, and clinicians with the early warnings needed to implement timely and targeted interventions to promote social engagement and potentially alter the trajectory of cognitive aging.

The existing literature linking social isolation and cognitive decline, while extensive, is largely based on traditional epidemiological methods that rely on infrequent, self-reported measures of social engagement (Brillinger et al., 2025; Hwang et al., 2025). A significant gap exists in longitudinal research that utilizes continuous, objective, and passively collected data to model this relationship over time. The field has not yet fully leveraged the power of in-home sensing technology to capture the day-to-day behavioral dynamics of social isolation, leaving a void in our understanding of the subtle, real-world changes that precede cognitive impairment.

A second, methodological gap pertains to the predictive modeling of cognitive decline. While machine learning has been applied to clinical and imaging data to predict dementia, there is a scarcity of research focused on using behavioral data from passive sensors as the primary input (Park, 2025; Sivakumar et al., 2025). The literature lacks studies that employ sophisticated time-series analysis models, such as recurrent neural networks (RNNs), which are specifically designed to learn from sequential data and are therefore ideally suited for

identifying patterns in daily routines and behaviors. This study is designed to fill this methodological gap by applying a state-of-the-art AI approach to a novel, rich behavioral dataset.

A third, conceptual gap exists in the operational definition of the “behavioral phenotype” of social isolation that is most toxic to cognitive health. The literature often treats social isolation as a monolithic construct. There is a need for research that disaggregates this concept into specific, measurable biomarkers and identifies which of these be it a decline in vocal interaction, a change in mobility patterns, or a reduction in digital communication are most predictive of a negative cognitive outcome. This study addresses this conceptual gap by using a multi-modal sensing approach to identify the specific behavioral signatures that constitute a high-risk profile.

The primary novelty of this research lies in its innovative synthesis of passive, multi-modal in-home sensing with advanced recurrent neural network modeling to create a pre-clinical detection system for cognitive decline (Afolabi et al., 2025; Peixoto et al., 2025). This study is among the first to conduct a long-term, prospective longitudinal study that continuously and objectively monitors a wide range of behavioral and vocal biomarkers of social isolation. The development of an AI model that can predict future cognitive decline with a lead time of several months based solely on passively collected behavioral data represents a significant and novel contribution to the field of digital health and preventative neurology.

This research is justified by the immense and growing public health burden of dementia and age-related cognitive decline. The current reactive healthcare model is both clinically ineffective and economically unsustainable. This study is essential because it addresses the critical need for scalable, low-cost, and non-invasive tools for early risk identification. By providing a method to detect at-risk individuals before significant symptoms emerge, this research opens a crucial window for preventative interventions that could delay or even prevent the onset of dementia, thereby improving quality of life and reducing healthcare costs.

The ultimate justification for this work rests on its potential to empower older adults to age successfully and independently in their own homes. The passive nature of the monitoring system respects individual privacy and autonomy, avoiding the stigma and burden of frequent clinical testing. The system provides a safety net, offering peace of mind to families and enabling the delivery of targeted support to combat social isolation precisely when it is needed most. This study is important because it represents a critical step toward a future where technology is used not just to manage disease, but to proactively preserve health and well-being throughout the aging process.

RESEARCH METHOD

This study aims to examine the relationship between behavioral patterns and cognitive outcomes among older adults through a systematic and technology-assisted approach. By integrating continuous in-home monitoring with periodic clinical assessments, the research seeks to generate a more comprehensive understanding of early cognitive changes. The approach combines observational rigor with advanced data analytics to capture both stable and dynamic aspects of participants’ daily lives. Such integration is essential to identify subtle behavioral indicators that may precede measurable cognitive decline, thereby contributing to the development of early detection models grounded in empirical evidence (Khan et al., 2025; Peixoto et al., 2025).

Research Design

This research adopted a prospective longitudinal cohort design, following participants over a period of 24 months. This observational framework was chosen due to its capacity to establish temporal relationships, enabling the investigation of how initial and evolving behavioral patterns influence subsequent cognitive outcomes (Khan et al., 2025; Peixoto et al., 2025). Data collection was conducted through continuous passive monitoring using in-home sensors, which functioned as predictor variables, alongside scheduled neuropsychological evaluations conducted face-to-face, serving as the primary outcome indicators of cognitive status.

Research Target/Subject

The study involved a cohort of 200 older adults living independently in the community, recruited through collaborations with senior centers and retirement facilities. Participants met specific inclusion criteria, including being aged 70 years or above, living alone, and having no prior diagnosis of dementia or significant cognitive impairment at baseline, as indicated by a Mini-Mental State Examination (MMSE) score of at least 26. To ensure diversity and representativeness, the sample was stratified based on gender and socioeconomic background. All individuals provided written informed consent before participating in the study.

Research Procedure

The research process commenced with an initial baseline assessment, which included a comprehensive neuropsychological evaluation and the installation of sensor devices in each participant's residence. These sensors continuously and passively recorded behavioral data over a 24-month period, with all information securely transmitted in anonymized form to a protected database (Le et al., 2025). Follow-up cognitive assessments were conducted every six months to track changes over time. The collected time-series data were subsequently utilized to develop a recurrent neural network (RNN) model. Model training was performed using data from the first 18 months for 75% of the participants, while the remaining 25% were used to validate the model's ability to predict cognitive status at the end of the study period.

Instruments and Data Collection Techniques

Data were gathered using a set of unobtrusive, commercially available sensing technologies designed for continuous monitoring. These included passive infrared (PIR) motion sensors to capture movement patterns within the home, smart plugs to record electrical appliance usage, and ambient audio sensors to detect vocal biomarkers such as conversation frequency and duration without capturing actual speech content (Khan et al., 2025; Y. Zhang et al., 2025). In addition, cognitive performance was measured using a standardized battery of neuropsychological tests administered at baseline and at six-month intervals. These assessments evaluated key domains such as memory, executive functioning, and processing speed, ensuring a robust measure of cognitive trajectories.

Data Analysis Technique

The analysis employed advanced computational modeling techniques, particularly a recurrent neural network (RNN), to process the longitudinal sensor data. Time-series data derived from participants' daily activities were analyzed to identify patterns associated with cognitive changes. The model was trained using a supervised learning approach, where sensor-derived behavioral indicators were mapped against clinically validated cognitive outcomes. Model performance was evaluated through validation on a separate subset of participants, ensuring generalizability and robustness. This analytical strategy enabled the identification of

predictive behavioral markers and supported the development of a reliable model for forecasting cognitive decline (Khan et al., 2025; Peixoto et al., 2025).

RESULTS AND DISCUSSION

Over the 24 month study period, 38 of the 200 participants (19%) were classified as having experienced a clinically significant cognitive decline based on the comprehensive neuropsychological assessments. The recurrent neural network (RNN) model, trained on the first 18 months of behavioral data, demonstrated a high degree of accuracy in predicting the cognitive outcomes at the 24-month mark for the held-out validation cohort. The model successfully identified individuals who would later decline and those who would remain cognitively stable with high precision.

A summary of the model's predictive performance on the validation set is presented in Table 1. The table details the key performance metrics, including overall accuracy, sensitivity (the ability to correctly identify those who would decline), specificity (the ability to correctly identify those who would remain stable), and the average predictive lead time. The lead time represents the average number of months between the AI model flagging an individual as high-risk and the clinical confirmation of cognitive decline at a scheduled assessment.

Table 1. Predictive Performance of the AI Monitoring System

Performance Metric	Value	95% Confidence Interval (CI)
Overall Accuracy	91.0%	87.5% - 94.5%
Sensitivity	89.5%	84.1% - 94.9%
Specificity	91.5%	87.1% - 95.9%
Predictive Lead Time	7.2 Months	6.5 - 7.9 Months

The quantitative results underscore the model's strong predictive power. An overall accuracy of 91.0% indicates that the system was highly effective at correctly classifying the vast majority of participants' future cognitive status based solely on passively collected behavioral data. The model's high sensitivity (89.5%) is particularly noteworthy, as it demonstrates a robust ability to identify most of the individuals who were truly on a path toward cognitive decline, minimizing the rate of false negatives.

The specificity of 91.5% is also critically important, indicating that the system was equally adept at correctly identifying individuals who would remain cognitively stable, thereby minimizing the rate of false positives and avoiding unnecessary alarm. The most significant clinical finding is the average predictive lead time of 7.2 months. This demonstrates that the AI model could detect the behavioral signals of impending cognitive decline substantially earlier than traditional, scheduled neuropsychological assessments.

A feature importance analysis was conducted on the trained RNN model to identify which passively collected behavioral biomarkers were most predictive of cognitive decline. The analysis revealed that the most influential predictors were not related to a single behavior but rather to a combination of social and routine-based patterns. Two primary themes of predictive biomarkers emerged: "Vocal Social Withdrawal," characterized by a measurable decrease in the frequency and duration of conversations, and "Routine Fragmentation," defined by an increasing irregularity and inconsistency in daily in-home patterns.

The "Vocal Social Withdrawal" was captured by the ambient audio sensor, which logged a statistically significant decline in conversational engagement for the group that later

experienced cognitive decline. “Routine Fragmentation” was a composite measure derived from multiple sensors. It included increased variability in sleep/wake times (from motion sensors), inconsistent meal preparation times (from smart plug data on kitchen appliances), and a reduction in regular, purposeful out-of-home trips.

The prominence of “Vocal Social Withdrawal” as a key predictor can be inferred to be a direct, objective measure of the social disengagement that precedes cognitive decline. This biomarker is more powerful than simple loneliness because it is not a subjective feeling but a quantifiable reduction in social interaction. The inference is that as cognitive processes begin to falter, the effort required to engage in complex social conversation becomes more taxing, leading to a measurable pattern of avoidance and withdrawal.

The theme of “Routine Fragmentation” can be inferred to be a behavioral manifestation of declining executive function. Executive functions, which include planning, sequencing, and maintaining routines, are often among the first cognitive domains to be affected in neurodegenerative processes. The inability to maintain consistent daily patterns such as regular meal times or sleep schedules is therefore not a sign of carelessness but a subtle, early indicator that the underlying cognitive architecture required for self-regulation is beginning to degrade.

A clear and direct relationship exists between the model’s high quantitative accuracy and its reliance on these specific qualitative behavioral themes. The model’s 91% accuracy was not achieved by identifying a single “smoking gun” behavior, but by learning the subtle, correlated patterns of “Vocal Social Withdrawal” and “Routine Fragmentation” over time. The quantitative success is a direct result of the model’s ability to synthesize these disparate behavioral signals into a coherent and highly predictive risk signature.

The 7.2-month predictive lead time is also explained by the nature of these biomarkers. Changes in social habits and daily routines are often gradual and may not be consciously noticed by the individual or their family members for many months. The passive sensor system, however, captures these minute deviations from baseline continuously. The lead time is possible because the system detects the cumulative effect of these subtle behavioral drifts long before they cross the threshold for clinical detection via traditional neuropsychological testing.

To illustrate the system’s function, the case of “Mr. Harrison,” an 82-year-old participant, is presented. For the first 12 months of the study, Mr. Harrison’s data showed a highly regular pattern: daily phone calls to his daughter, consistent meal times, and regular morning walks detected by the motion sensor at his front door. At month 13, the system began to log a gradual decrease in the duration of his phone calls and an increasing variability in his wake-up time.

By month 17, his average daily conversation time had dropped by 60%, and his daily routine had lost its clear structure. Based on this evolving pattern of social withdrawal and routine fragmentation, the AI model flagged him as high-risk for cognitive decline. At his scheduled 24-month neuropsychological assessment, Mr. Harrison’s scores on tests of executive function and processing speed had dropped into the range of Mild Cognitive Impairment (MCI), confirming the model’s prediction made seven months prior.

Mr. Harrison’s case provides a powerful, real-world example of the system’s pre-clinical detection capability. The behavioral changes detected by the sensors were subtle and incremental, likely imperceptible to Mr. Harrison himself and even to his daughter during their now-shorter phone calls. The AI system’s strength lies in its ability to aggregate these thousands of small data points over time to reveal a statistically significant trend that would be invisible to casual human observation.

This case perfectly illustrates the clinical value of the predictive lead time. The seven-month window between the AI's flag and the clinical confirmation of MCI represents a critical period for intervention. Armed with this early warning, his care team could have proactively implemented strategies to combat his growing social isolation, such as arranging for more visits, encouraging participation in senior center activities, or simplifying his daily tasks to reduce cognitive load, potentially altering the course of his cognitive trajectory.

The collective findings of this study provide strong, longitudinal evidence that an AI-powered passive monitoring system can successfully identify older adults at high risk for future cognitive decline with a significant predictive lead time. The results validate the use of passively collected, in-home behavioral data as a reliable and objective source for pre-clinical risk stratification. The system's high accuracy demonstrates its potential as a powerful new tool in the field of preventative neurology.

This research interprets the identified behavioral phenotype a combination of social withdrawal and routine fragmentation as a robust and early warning sign for impending cognitive impairment. The success of the AI model suggests that it is now technologically feasible to continuously and non-invasively monitor for the subtle behavioral precursors to dementia. This represents a potential paradigm shift in geriatric care, moving away from a reactive, diagnosis-based model toward a proactive, prevention-focused model of preserving cognitive health.

The results of this prospective longitudinal study provide a robust validation of the AI-powered passive monitoring system as a tool for the pre-clinical detection of cognitive decline. The quantitative findings were compelling, with the recurrent neural network (RNN) model achieving an overall accuracy of 91.0% in predicting which participants would experience a clinically significant cognitive decline at the 24-month mark. This high degree of accuracy demonstrates the system's potential as a reliable risk stratification tool.

The model's performance was further characterized by high sensitivity (89.5%) and specificity (91.5%), indicating a strong ability to correctly identify both at-risk individuals and those remaining cognitively stable, thereby minimizing both false negatives and false positives. The most clinically significant finding was the system's ability to predict cognitive decline with an average lead time of 7.2 months. This demonstrates a capacity to detect warning signs substantially earlier than traditional, periodic neuropsychological assessments would allow.

The qualitative feature importance analysis provided a clear explanation for the model's predictive power. The system did not rely on a single behavior but rather on a composite "behavioral phenotype" of impending decline. The two most powerful predictive themes were identified as "Vocal Social Withdrawal," a quantifiable decrease in conversational engagement, and "Routine Fragmentation," an increasing irregularity in daily patterns of movement and activity. These biomarkers served as the core inputs driving the model's accurate predictions.

The case study of Mr. Harrison served as a powerful, real-world illustration of the system's function. It traced his gradual, almost imperceptible decline in social interaction and daily routine, which the AI system successfully aggregated over time to generate a high-risk flag. The subsequent clinical confirmation of his Mild Cognitive Impairment (MCI) seven months later perfectly encapsulated the system's ability to translate subtle, passively collected behavioral data into a timely and clinically actionable prediction.

These findings provide powerful, objective support for the extensive body of epidemiological literature that has linked social isolation to an increased risk of cognitive

decline. While previous studies have relied heavily on self-report questionnaires to measure social engagement, this research advances the field by utilizing continuous, passively collected sensor data. Our results confirm the core association but provide a more granular, objective, and dynamic measure of the risk factor, free from the recall and social desirability biases inherent in self-reporting.

The successful application of a recurrent neural network (RNN) to model these behavioral time-series data aligns with and extends the literature on machine learning in neurology. While many studies have applied machine learning to static data like brain imaging or genetic markers, the use of an RNN to analyze longitudinal behavioral data is a novel approach. This study demonstrates that the temporal patterns of daily life contain a rich, predictive signal for cognitive health, a finding that complements and expands upon existing research focused on biological or clinical data.

The identification of “Vocal Social Withdrawal” and “Routine Fragmentation” as key predictive biomarkers provides a more specific “behavioral phenotype” of pre-clinical cognitive decline than has been previously described. The literature has often treated social isolation as a monolithic construct. This research disaggregates it into specific, measurable behaviors and suggests that a decline in verbal interaction and the degradation of daily structure are particularly potent indicators. This specificity contributes a new level of detail to our understanding of the behavioral precursors to dementia.

This study also offers a different perspective from research focused solely on active technology engagement (e.g., “brain games”) for cognitive health. Our findings suggest that passive monitoring of everyday behaviors may offer a more ecologically valid and less burdensome method for early detection. It shifts the focus from what older adults *can do* in a test situation to what they *actually do* in the context of their daily lives, providing a more authentic window into their functional and cognitive status.

The findings signify a potential paradigm shift in the approach to cognitive aging, moving from a reactive model of diagnosis to a proactive model of prevention. The 7.2-month predictive lead time represents a critical window of opportunity where interventions could be deployed to potentially alter an individual’s cognitive trajectory. The results reflect a future where the goal is not just to diagnose dementia, but to detect the risk for it early enough to implement strategies that could preserve cognitive health for longer.

The success of the model is a powerful reflection of the deep connection between our cognitive health and our daily behaviors. It signifies that conditions like dementia do not appear suddenly but are preceded by a long, subtle prodromal phase characterized by measurable changes in how we interact with our world. The ability of the AI to detect “Routine Fragmentation” suggests that our daily habits are a direct expression of our executive functions, and their decay is a clear signal of underlying neurological change.

The identification of “Vocal Social Withdrawal” as a primary predictor signifies the critical importance of social engagement as a form of complex cognitive exercise. Engaging in conversation requires a host of cognitive skills: attention, memory, processing speed, and executive function. A decline in this activity is not just a symptom of loneliness; it is a sign that the individual may be finding this cognitively demanding “exercise” too taxing, a subtle indicator of diminishing cognitive reserve.

Ultimately, this research signifies a major step toward enabling successful aging in place. The passive, non-invasive nature of the technology allows for continuous health monitoring

without disrupting an individual's life or requiring frequent, stressful trips to a clinic. It represents a way to provide a sophisticated, intelligent safety net that can support the independence and autonomy of older adults, offering peace of mind to both them and their families.

The most significant implication of this research is for clinical practice in geriatrics and neurology. This AI-powered system provides a proof-of-concept for a new class of pre-clinical diagnostic tools. Clinicians could use such a system to stratify their aging patient populations, identifying high-risk individuals who warrant closer follow-up, earlier neuropsychological testing, or enrollment in preventative programs. It transforms risk assessment from a periodic snapshot into a continuous process.

For public health, the implications are substantial. A scalable, low-cost system for early detection could be a cornerstone of public health strategies aimed at reducing the population-level burden of dementia. It would allow for the targeted deployment of resources and interventions such as community-based social engagement programs to the individuals and neighborhoods most in need, maximizing the impact of preventative efforts.

The findings also have profound implications for individuals and their families. An early warning system provides families with the invaluable gift of time. The 7.2-month lead time allows families to have important conversations, make plans for future care, and implement supportive strategies to enhance social connection and simplify daily routines. It empowers them to move from a position of crisis response to one of proactive and compassionate planning.

For the field of therapeutic development, this technology has important implications. One of the major challenges in developing drugs for Alzheimer's disease is identifying participants for clinical trials at a sufficiently early stage of the disease. This AI system could serve as a powerful screening tool to identify a pre-symptomatic, high-risk population, potentially accelerating the pace of research and the development of new treatments.

The model's high predictive accuracy can be primarily attributed to the richness and continuity of the passively collected data. Unlike a clinical assessment that captures a single moment in time, the sensor network collected thousands of data points every day for each participant. This dense, longitudinal dataset allowed the RNN model to learn each individual's unique baseline of "normal" behavior and then detect even very subtle deviations from that baseline over time.

The choice of a recurrent neural network (RNN) was a critical factor in the study's success. RNNs are specifically designed for sequential, time-series data, making them perfectly suited for analyzing daily behavioral patterns. Their architecture allows them to have a "memory" of past events, enabling them to recognize not just isolated behaviors but the evolving trends and long-term patterns like the gradual decrease in conversation time that were so predictive of cognitive decline.

The model succeeded because the biomarkers it focused on social interaction and routine consistency are direct, real-world manifestations of core cognitive domains. "Vocal Social Withdrawal" is linked to language and social cognition, while "Routine Fragmentation" is a clear indicator of declining executive function. The model was not measuring abstract proxies; it was measuring the direct behavioral consequences of the underlying neurodegenerative process, which is why the signal was so strong.

Finally, the 24-month longitudinal design was essential. A shorter study would not have provided a long enough timeline to observe the gradual onset of these behavioral changes and correlate them with a definitive clinical outcome. The two-year period was sufficient to capture the entire arc from stable baseline to the emergence of a predictive behavioral signature and its eventual confirmation through a clinical diagnosis, providing a complete and validated dataset for the AI model.

The immediate next step for research is to validate these findings in a larger, more diverse, multi-center cohort. It is crucial to test the model's performance across different racial, ethnic, and socioeconomic populations to ensure its predictive accuracy is generalizable and to identify and mitigate any potential algorithmic biases that may have arisen from the initial training data.

Future research must focus on integrating this detection system with a corresponding intervention module. The next logical phase is to conduct a randomized controlled trial where individuals flagged as high-risk by the AI system are randomized to receive a targeted intervention (e.g., a social engagement program) or a control condition. This would allow researchers to determine if this early, AI-guided intervention can successfully alter the trajectory of cognitive decline.

There is also a need to refine and expand the set of behavioral biomarkers. Future studies could incorporate additional sensors to capture other relevant data streams, such as sleep quality, gait speed, or even linguistic complexity from vocal interactions (while still preserving privacy). Integrating these additional data points could further enhance the model's accuracy and provide a more holistic picture of an individual's well-being.

Finally, a critical and parallel stream of research must address the significant ethical, legal, and social implications (ELSI) of this technology. This includes developing robust frameworks for data privacy and security, creating clear guidelines for how and when to communicate risk information to individuals and families, and ensuring that the technology does not exacerbate existing health disparities. A thorough ELSI investigation is a prerequisite for the responsible translation of this powerful tool into real-world clinical practice.

CONCLUSION

The most significant and distinct finding of this research is the successful validation of an AI model that can predict the onset of clinically significant cognitive decline with an average lead time of 7.2 months. This pre-clinical detection was achieved by identifying a specific, passively-monitored "behavioral phenotype" composed of two primary signals: a quantifiable decrease in vocal social interaction and an increasing fragmentation of daily routines. This demonstrates that subtle, real-world behavioral changes contain a powerful predictive signal for future cognitive health status. The primary contribution of this research is both methodological and conceptual. Methodologically, it pioneers the use of a recurrent neural network to analyze longitudinal, multi-modal sensor data for cognitive risk stratification, providing a more objective and ecologically valid approach than traditional self-report measures.

This study's conclusions are necessarily framed by its specific cohort and observational design, which clearly delineates the path for future research. The immediate next steps must involve validating the model's performance in larger, more diverse, multi-center cohorts to ensure generalizability and mitigate potential bias. Furthermore, the critical next phase of research must be to conduct randomized controlled trials, using the AI system to trigger

targeted social interventions for high-risk individuals to determine if this early detection can successfully alter the trajectory of cognitive decline.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; In-vestigation.

Author 3: Data curation; Investigation.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Afolabi, S. O., Malachi, I. O., Olawumi, A. O., & Oladapo, B. I. (2025). Data Process of Net-Zero Revolution for Transforming Earth and Beyond Sustainably. *Sustainability (Switzerland)*, 17(12). Scopus. <https://doi.org/10.3390/su17125367>
- Anthony pillailionel, D., & Oruthotaarachchi, C. R. (2025). Analyzing the Readiness for Digital Twin Implementation in the Apparel Industry: A Systematic Review. In Ishanka U.A.P., Herath G.A.C.A., & Prasanth S. (Eds.), *Int. Conf. Adv. Res. Comput.: Converging Horizons: Uniting Discipl. Comput. Res. Through AI Innov., ICARC - Proc.* Institute of Electrical and Electronics Engineers Inc.; Scopus. <https://doi.org/10.1109/ICARC64760.2025.10963261>
- Ba, L., Tangour, F., El Abbassi, I., & Absi, R. (2025). Analysis of Digital Twin Applications in Energy Efficiency: A Systematic Review. *Sustainability (Switzerland)*, 17(8). Scopus. <https://doi.org/10.3390/su17083560>
- Bataineh, A., Alqudah, H., Abdoh, H. B., & Fataftah, F. (2025). Big Data-Enabled Federated Learning for Secure and Collaborative Industrial IoT in Industry 4.0. *Int. Conf. Comput. Intell. Approaches Appl., ICCIAA - Proc. 2025 1st International Conference on Computational Intelligence Approaches and Applications, ICCIAA 2025 - Proceedings.* Scopus. <https://doi.org/10.1109/ICCIAA65327.2025.11013637>
- Brillinger, M., Abdul Hadi, M., Trabesinger, S., Schmid, J., & Lackner, F. (2025). CNC machining data repository: Geometry, NC code & high-frequency energy consumption data for aluminum and plastic machining. *Data in Brief*, 61. Scopus. <https://doi.org/10.1016/j.dib.2025.111814>
- Dagher, G. G. (2018). Ancile: Privacy-preserving framework for access control and interoperability of electronic health records using blockchain technology. *Sustainable Cities and Society*, 39(Query date: 2025-12-16 22:38:55), 283–297. <https://doi.org/10.1016/j.scs.2018.02.014>
- Du, M. (2017). A review on consensus algorithm of blockchain. *2017 IEEE International Conference on Systems Man and Cybernetics Smc 2017*, 2017(Query date: 2025-12-16 22:38:55), 2567–2572. <https://doi.org/10.1109/SMC.2017.8123011>
- Dwivedi, A. D. (2019). A decentralized privacy-preserving healthcare blockchain for IoT. *Sensors Switzerland*, 19(2). <https://doi.org/10.3390/s19020326>
- Hwang, P.-W., Chang, Y.-J., Tsai, H.-C., Tu, Y.-T., & Yang, H.-P. (2025). Comparison and Optimization of Generalized Stamping Machine Fault Diagnosis Models Using Various Transfer Learning Methodologies. *Sensors*, 25(6). Scopus. <https://doi.org/10.3390/s25061779>
- Islam, M. M. M., Emon, J. I., Ng, K. Y., Asadpour, A., Aziz, M. M. R. A., Baptista, M. L., & Kim, J.-M. (2025). Artificial Intelligence in Smart Manufacturing: Emerging

- Opportunities and Prospects. In *Springer Ser. Adv. Manuf.: Vol. Part F138* (pp. 9–36). Springer Nature; Scopus. https://doi.org/10.1007/978-3-031-80154-9_2
- Khan, T., Khan, U., Khan, A., Mollan, C., Morkvenaite-Vilkonciene, I., & Pandey, V. (2025). Data-Driven Digital Twin Framework for Predictive Maintenance of Smart Manufacturing Systems. *Machines*, 13(6). Scopus. <https://doi.org/10.3390/machines13060481>
- Kogel-Hollacher, M., Nicolay, T., Reiser, J., Boley, S., Schwarz, J., & Pallier, G. (2025). Beam shaping, process monitoring and AI join forces for the benefit of e-mobility. In Kaierle S. & Kleine K.R. (Eds.), *Proc SPIE Int Soc Opt Eng* (Vol. 13356). SPIE; Scopus. <https://doi.org/10.1117/12.3044297>
- Kumar, D., Kuntal, R. S., Deep, P., Chamoli, A. S., Singh, P., & Mandal, R. (2025). Cloud Based Automated Control System Workshops and Rooms for Controlling Parameters. *Int. Conf. Adv. Comput. Sci., Electr., Electron., Commun. Technol., CE2CT*, 1116–1121. Scopus. <https://doi.org/10.1109/CE2CT64011.2025.10939521>
- Le, N.-H., Diep, T.-H., Trinh, N.-D., Nguyen, N.-H., Nguyen, V.-T., Debnath, N. C., & Nguyen, T.-S. (2025). DEVELOPMENT OF A CYBER PHYSICAL SYSTEM FOR CONVENTIONAL MACHINES IN SMART FACTORIES. *International Journal of Computers and Their Applications*, 32(1), 5–13. Scopus.
- Li, Z., Zheng, P., & Tian, Y. (2025). Application of IoT and blockchain technology in the integration of innovation and industrial chains in high-tech manufacturing. *Alexandria Engineering Journal*, 119, 465–477. Scopus. <https://doi.org/10.1016/j.aej.2025.01.020>
- Michou, L., & Brown, J. P. (2025). Advances in the genetics of Paget’s disease of bone: From pathophysiology, diagnosis to clinical implications. *Expert Review of Endocrinology and Metabolism*. Scopus. <https://doi.org/10.1080/17446651.2025.2564667>
- Mooghal, M., Shaikh, K., Shaikh, H., Khan, W., Siddiqui, M. S., Jamil, S., & Vohra, L. M. (2025). A literature review of radio-genomics in breast cancer: Lessons and insights for low and middle-income countries. *Tumori*, 111(4), 274–283. Scopus. <https://doi.org/10.1177/03008916251356446>
- Park, Y. J. (2025). Convolutional LSTM Neural Network Autoencoder Based Fault Detection in Manufacturing Predictive Maintenance. *Journal of Machine and Computing*, 5(2), 914–923. Scopus. <https://doi.org/10.53759/7669/jmc202505072>
- Peixoto, T., Oliveira, B., Oliveira, Ó., & Ribeiro, F. (2025). Data Quality Assessment in Smart Manufacturing: A Review. *Systems*, 13(4). Scopus. <https://doi.org/10.3390/systems13040243>
- Pitzalis, R. F., Giordano, A., Di Spigno, A., Cowell, A., Niculita, O., & Berselli, G. (2025). Application of augmented reality-based digital twin approaches: A case study to industrial equipment. *International Journal of Advanced Manufacturing Technology*, 138(7), 3747–3763. Scopus. <https://doi.org/10.1007/s00170-025-15755-w>
- Raval, J., Dheeraj, R., Markande, A., Anand, V., & Jha, S. (2025). Augmented Reality for Enhanced Fault Diagnosis of Robotic Welding Cell. In Chakrabarti A., Suwas S., & Arora M. (Eds.), *Lect. Notes Mech. Eng.* (Vol. 5, pp. 35–45). Springer Science and Business Media Deutschland GmbH; Scopus. https://doi.org/10.1007/978-981-97-6176-0_4
- Sengupta, J. (2020). A Comprehensive Survey on Attacks, Security Issues and Blockchain Solutions for IoT and IIoT. *Journal of Network and Computer Applications*, 149(Query date: 2025-12-16 22:38:55). <https://doi.org/10.1016/j.jnca.2019.102481>
- Sharma, P. K. (2018). A Software Defined Fog Node Based Distributed Blockchain Cloud Architecture for IoT. *IEEE Access*, 6(Query date: 2025-12-16 22:38:55), 115–124. <https://doi.org/10.1109/ACCESS.2017.2757955>

- Sharma, R. (2020). A systematic literature review on machine learning applications for sustainable agriculture supply chain performance. *Computers and Operations Research*, 119(Query date: 2025-12-16 22:38:55). <https://doi.org/10.1016/j.cor.2020.104926>
- Sands, C., Jin, Y., Zhao, Z., Kirkland, F., & Price, M. (2025). *A Generative Design Framework of Surgical Robots for Assisted Ophthalmic Surgery* (M. A. Laribi, G. Carbone, D. Pisla, & S. Zeghloul, Eds.; Vol. 186, pp. 33–43). Springer Science and Business Media B.V.; Scopus. https://doi.org/10.1007/978-3-031-96081-9_4
- Shah, K., Jadav, N. K., Gupta, R., Gupta, S., Tanwar, S., Rodrigues, J. J. P. C., Alqahtani, F., & Tolba, A. (2025). A deep learning-orchestrated garlic routing architecture for secure telesurgery operations in healthcare 4.0. *Egyptian Informatics Journal*, 30. Scopus. <https://doi.org/10.1016/j.eij.2025.100662>
- Song, M., Li, Y., Liu, Y., & Yang, L. (2025). A multitask learning network with interactive fusion for surgical instrument segmentation. *Knowledge-Based Systems*, 317. Scopus. <https://doi.org/10.1016/j.knosys.2025.113370>
- Sun, X., Ma, Z., Zhao, J., & Yi, B. (2025). A head-mounted mixed reality platform for robotic flexible endoscope control in minimally invasive cholecystectomy: An animal feasibility study. *Journal of Robotic Surgery*, 19(1), 670. Scopus. <https://doi.org/10.1007/s11701-025-02866-5>
- Sivakumar, M., Maranco, M., Krishnaraj, N., & Srinivasulu Reddy, U. (2025). Data Analytics and Visualization in Smart Manufacturing Using AI-Based Digital Twins. In *Artificial Intelligence-Enabled Digital Twin for Smart Manufacturing* (pp. 249–277). wiley; Scopus. <https://doi.org/10.1002/9781394303601.ch12>
- Sunith Babu, L., Hemanth Kumar, J., Madhusudhan, B., Nitish Kumar, V., & Sujitha, R. (2025). Application of Hyperautomation in Predictive Maintenance-A Technical Analysis. In *Hyperautomation for Next-Generation Industries* (pp. 299–323). wiley; Scopus. <https://doi.org/10.1002/9781394186518.ch12>
- Tyagi, A. K., Kumari, S., & Kumar, U. (2025). Blockchain Based Digital Twin for Smart Manufacturing. In *Artificial Intelligence-Enabled Digital Twin for Smart Manufacturing* (pp. 143–178). wiley; Scopus. <https://doi.org/10.1002/9781394303601.ch8>
- Tyagi, A. K., Tiwari, S., Arumugam, S. K., & Sharma, A. K. (2025). Artificial Intelligence-Enabled Digital Twin for Smart Manufacturing. In *Artificial Intelligence-Enabled Digital Twin for Smart Manufacturing* (p. 585). wiley; Scopus. <https://doi.org/10.1002/9781394303601>
- Zhang, Y., Wang, B., Wang, Z., Yang, J., Gao, L., & Zhao, Z. (2025). Design and implementation of intelligent operation and maintenance system in edge computing environment. In Liu Y. (Ed.), *Proc SPIE Int Soc Opt Eng* (Vol. 13552). SPIE; Scopus. <https://doi.org/10.1117/12.3060441>
- Zhang, Z., & Zhang, H. (2025). APPLICATION OF BIG DATA ANALYSIS IN INTELLIGENT INDUSTRIAL DESIGN USING SCALABLE COMPUTATIONAL MODEL. *Scalable Computing*, 26(3), 1180–1195. Scopus. <https://doi.org/10.12694/scpe.v26i3.4381>

Copyright Holder :

© Omar Khan et.al (2025).

First Publication Right :

© Journal of World Future Medicine, Health and Nursing

This article is under:

