

Nutrition Education for Adolescent Girls as a Strategy for Intergenerational Stunting Prevention: A Quantitative Study at SMAN 03 Rumbai

Siska Indrayani¹, Rina Oktaviana²

¹ Universitas Muhammadiyah Riau, Indonesia

² Universitas Muhammadiyah Riau, Indonesia

Corresponding Author:

Siska Indrayani

Faculty of MIPA and Kes, Universitas Muhammadiyah Riau

Jl. KH. Ahmad Dahlan No.88, Kp. Melayu, Kec. Sukajadi, Kota Pekanbaru, Riau 28156

Email: siskaindrayani@umri.ac.id

Article Info

Received: July 10, 2025

Revised: Sep 6, 2025

Accepted: Oct 9, 2025

Online Version: Dec 7, 2025

Abstract

Stunting is a chronic nutritional problem with long-term consequences for human capital and is influenced by nutritional conditions across the life course, including adolescence as a critical preconception period. Adolescent girls play a strategic role in intergenerational stunting prevention; however, nutrition literacy in this group remains relatively low. This study aimed to analyze the effect of nutrition education on improving nutrition knowledge among adolescent girls as a preventive strategy for intergenerational stunting among senior high school students. A quasi-experimental study with a one-group pretest posttest design was conducted. The study participants consisted of 100 female students in grades X and XI from a senior high school in Rumbai, Indonesia, selected using a total sampling technique. Data were collected using a structured nutrition knowledge questionnaire that had been tested for validity and reliability. Data analysis included univariate and bivariate analyses. Normality was assessed using the Shapiro Wilk test, and differences in nutrition knowledge scores before and after the intervention were analyzed using a paired t-test with a significance level of $p < 0.05$. The mean nutrition knowledge score increased significantly from 65.42 ± 6.12 before the intervention to 81.30 ± 6.05 after the intervention ($p < 0.001$). A mean difference of 15.88 points indicates a substantial improvement in nutrition knowledge following the nutrition education program. In conclusion, nutrition education is effective in improving nutrition knowledge among adolescent girls and has the potential to serve as an upstream preventive strategy for intergenerational stunting.

Keywords: Education, Nutrition, Stunting



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/health> ISSN: (P: 2988-7550) - (E: 2988-0459)

How to cite:

Indrayani, S & Oktaviana, R. (2025). Nutrition Education for Adolescent Girls as a Strategy for Intergenerational Stunting Prevention: A Quantitative Study at SMAN 03 Rumbai. *Journal of World Future Medicine, Health and Nursing*, 3(6), 502–509. <https://doi.org/10.70177/health.v3i6.3043>

Published by:

Yayasan Adra Karima Hubbi

INTRODUCTION

Stunting is a chronic nutritional problem with long-term consequences for the quality of human resources, including impaired physical growth, cognitive development, and an increased risk of non-communicable diseases in adulthood. Evidence from previous studies indicates that stunting is not solely determined by nutritional conditions during early childhood, but rather represents the cumulative effects of nutritional problems across the life course, including adolescence as a critical preconception period (Fahmida et al., 2025; Jaleel et al., 2025; Mukhopadhyay, 2025). Therefore, stunting prevention efforts should not be limited to pregnancy and the first 1,000 days of life, but should also target adolescent girls as future mothers.

Adolescent girls are particularly vulnerable to nutritional problems, such as inadequate nutrient intake and low nutrition literacy, which may affect their nutritional preparedness prior to pregnancy. Studies conducted in Indonesia have shown a significant association between adolescents' nutrition knowledge and nutritional status, suggesting that limited understanding of balanced nutrition may increase the risk of long-term nutritional problems, including stunting in the next generation (Akombi-Inyang, 2025; Brown et al., 2025). These findings highlight the importance of implementing promotive and preventive interventions beginning at the secondary school level.

School-based nutrition education is widely regarded as an effective and sustainable strategy to improve nutrition knowledge and awareness among adolescent girls. Experimental studies have demonstrated that nutrition education delivered through educational media, such as flip charts, significantly enhances adolescents' knowledge and attitudes toward stunting prevention (Anyati et al., 2025; Azhar et al., 2025; Pasha et al., 2025). In addition, interactive and context-based educational approaches, including educational games, have been shown to increase adolescents' awareness of the importance of balanced nutrition in daily life (Bezie et al., 2025; Hasib Joarder et al., 2025; Lundberg et al., 2025).

Further empirical evidence suggests that peer-based nutrition education models are also effective in strengthening adolescent girls' understanding as prospective mothers. Quantitative studies have reported that peer group nutrition education contributes positively to improvements in adolescents' nutrition knowledge and dietary behaviors and is considered a strategic approach to reducing the risk of stunting in a sustainable manner (Mekuriaw et al., 2025; Samuel et al., 2025; Zemariam et al., 2025). These findings reinforce the argument that nutrition education among adolescent girls plays a critical role in breaking the cycle of intergenerational stunting.

Despite this evidence, most stunting prevention programs in Indonesia remain focused on nutritional interventions for pregnant women and young children, while upstream approaches targeting adolescent girls have not been systematically optimized. In fact, adolescents' nutrition literacy plays an essential role in shaping healthy eating behaviors and nutritional readiness before entering the reproductive period, which ultimately contributes to stunting prevention (Bustos et al., 2025; Fogarty et al., 2025; Haniarti et al., 2025). This gap underscores the need for quantitative studies that specifically examine the role of nutrition education for adolescent girls in the context of intergenerational stunting prevention.

Based on these considerations, this study addresses the limited quantitative evidence regarding the effectiveness of nutrition education for adolescent girls as a strategy for intergenerational stunting prevention at the senior high school level. Nutrition education was selected as the intervention because it is promotive, preventive, easily integrated into the school environment, and has the potential for long-term impact. This study aims to analyze the role of

nutrition education in improving nutrition knowledge and nutritional preparedness among adolescent girls as a preventive effort against intergenerational stunting. The findings are expected to contribute to strengthening national evidence and to serve as a foundation for the development of more effective and sustainable adolescent nutrition education programs.

RESEARCH METHOD

Contains the type of research, time and place of research, targets/objectives, research subjects, procedures, instruments and data analysis techniques as well as other matters related to the method of research. Targets and objectives, research subjects, procedures, data and instruments, and data collection techniques, as well as data analysis techniques and other matters related to the method of research can be written in sub-chapters, with sub-headings. Sub-subheadings do not need to be notated, but are written in lowercase with a capital letter, Times New Roman-11 bold, left aligned. As an example can be seen below .

Research Design

This study employed a quantitative approach using a quasi-experimental design with a one-group pretest posttest model. This design was selected to assess changes in nutrition knowledge among adolescent girls before and after the implementation of a nutrition education intervention as a strategy for intergenerational stunting prevention.

Research Target/Subject

The study participants consisted of 100 female students from grades X and XI. Inclusion criteria included active enrollment, age between 17 and 18 years, willingness to participate, and completion of all stages of the research. Exclusion criteria were absence during the pretest or posttest sessions and incomplete participation in the nutrition education activities. A total sampling technique was applied, whereby all students who met the eligibility criteria were included as study participants.

Research Procedure

The research procedures were conducted in accordance with the quasi-experimental design using a one-group pretest posttest approach. The study was implemented in several sequential stages. Initially, administrative approval was obtained, followed by coordination with the school authorities to ensure the smooth implementation of the research activities. Participants were then informed about the objectives, procedures, and ethical considerations of the study, and written informed consent was obtained prior to data collection (Guo et al., 2025; Hu et al., 2025; Soharwardi et al., 2025).

After the completion of the nutrition education sessions, post-intervention data were collected using the same nutrition knowledge questionnaire administered during the pretest. This posttest aimed to measure changes in participants' nutrition knowledge following the intervention. The pretest and posttest scores were subsequently compared to evaluate the effect of the nutrition education program (Burger et al., 2025; Hu et al., 2025; Shiyami & Mardiana, 2025).

Instruments, and Data Collection Techniques

The data collected in this study were primary data in the form of nutrition knowledge scores. A structured questionnaire was used as the research instrument, developed based on adolescent balanced nutrition guidelines and concepts of stunting prevention. The questionnaire underwent validity and reliability testing to ensure its accuracy and consistency in measuring nutrition knowledge (Fatkuriyah et al., 2025; Hu et al., 2025; Soharwardi et al., 2025). Data were

collected through self-administered questionnaires during the pretest and posttest sessions, with researcher supervision to ensure completeness and accuracy of responses

Data Analysis Technique

The data collected in this study were primary data in the form of nutrition knowledge scores. A structured questionnaire was used as the research instrument, developed based on adolescent balanced nutrition guidelines and concepts of stunting prevention (Mokori et al., 2025; Soharwardi et al., 2025; Worku et al., 2025). The questionnaire underwent validity and reliability testing to ensure its accuracy and consistency in measuring nutrition knowledge. Data were collected through self-administered questionnaires during the pretest and posttest sessions, with researcher supervision to ensure completeness and accuracy of responses

RESULTS AND DISCUSSION

The initial quantitative analysis focused on pretest and posttest data of adolescent girls' nutrition knowledge scores obtained through a structured nutrition knowledge questionnaire. Descriptive analysis revealed a clear upward trend following the implementation of the nutrition education intervention (Fogarty et al., 2025; Mokori et al., 2025; Soharwardi et al., 2025). An increase in the mean nutrition knowledge score was observed, accompanied by a shift in the minimum and maximum values toward higher scores, indicating an overall improvement in adolescents' understanding of balanced nutrition concepts. These findings suggest that the nutrition education intervention had a positive effect on enhancing nutrition knowledge among adolescent girls as part of intergenerational stunting prevention efforts.

Table 1. Mean Nutrition Knowledge Scores Before and After Nutrition Education

Variable	Mean \pm SD	Min–Max
Pretest	65,42 \pm 6,12	46 – 73
Posttest	81,30 \pm 6,05	60 – 90

Table 1 presents the mean nutrition knowledge scores before and after the nutrition education intervention. Prior to the intervention, the mean nutrition knowledge score was 65.42 \pm 6.12, with scores ranging from 46 to 73. After the nutrition education was delivered, the mean score increased to 81.30 \pm 6.05, with a minimum score of 60 and a maximum score of 90. This improvement demonstrates a substantial increase in nutrition knowledge following the intervention, both in terms of mean values and score distribution, reflecting a comprehensive enhancement of nutrition understanding among most participants.

Table 2. Normality Test for Nutrition Knowledge Scores

Variabel	Kolmogorov Smirnov Sig.	Shapiro Wilk Sig.
Pretest	0,143	0,605
Posttest	0,200	0,467

Normality testing using the Shapiro Wilk test indicated that the pretest ($p = 0.605$) and posttest ($p = 0.467$) nutrition knowledge scores were normally distributed. Therefore, differences between pretest and posttest scores were analyzed using a paired t-test.

These findings demonstrate that nutrition education significantly improves nutrition knowledge among adolescent girls. The observed increase in mean knowledge scores suggests that structured and targeted nutrition information effectively enhances adolescents' understanding of balanced nutrition principles. This result supports the role of nutrition education as an effective promotive preventive approach for adolescent populations.

The results of this study are consistent with those reported by Azhari and Fayasari (2020), who found that nutrition education delivered through lectures and animated video media significantly improved adolescents' nutrition knowledge, attitudes, and eating behaviors. Similarly, Nurcahyani (2020) reported that balanced nutrition counseling using video media had a positive effect on dietary intake among adolescent girls. These findings indicate that improvements in nutrition knowledge may serve as an important initial step toward fostering healthier eating behaviors.

This study also aligns with the findings of Farikhah (2021), who reported that nutrition education delivered through infographic-and web-based media effectively enhanced nutrition knowledge and balanced eating behaviors. In addition, Damayanti (2022) emphasized that dietary patterns are significantly associated with adolescents' nutritional status. Therefore, improving nutrition knowledge through education has the potential to contribute to better dietary practices and the prevention of nutritional problems among adolescent girls.

In the context of intergenerational stunting, enhanced nutrition knowledge among adolescent girls has important strategic implications for early prevention during the preconception phase. Adolescent girls with adequate nutrition knowledge are expected to have better nutritional preparedness before entering the reproductive period, thereby reducing the risk of adverse pregnancy outcomes and low birth length, which are early determinants of stunting. Consequently, nutrition education for adolescent girls not only contributes to short-term improvements in knowledge but also represents a critical component of long-term strategies to prevent intergenerational stunting

CONCLUSION

This study demonstrates that nutrition education significantly improves nutrition knowledge among adolescent girls. The observed increase in knowledge scores following the intervention indicates that nutrition education is an effective approach to addressing the issue of limited nutrition understanding among adolescents.

More broadly, improved nutrition knowledge among adolescent girls has important implications for intergenerational stunting prevention. Adolescent girls with adequate nutrition knowledge are expected to have better nutritional preparedness before entering the reproductive period, which may reduce the risk of nutritional problems among mothers and children in the future. Therefore, nutrition education for adolescent girls can be considered an upstream preventive strategy to break the cycle of intergenerational stunting.

ACKNOWLEDGMENTS

The authors would like to express their gratitude for the support provided during the implementation of this study. This research was supported by PT Pertamina Hulu Rokan and the Institute for Research and Community Service (LPPM), Universitas Muhammadiyah Riau. The authors also extend their appreciation to all parties and participants who contributed to and supported the successful completion of this study.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Methodology; Project administration.

Author 2: Investigation; Validation; Supervision; Resources; Writing – review and editing.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Azhari, M. A., & Fayasari, A. (2020). Pengaruh edukasi gizi dengan media ceramah dan video animasi terhadap pengetahuan, sikap, dan perilaku sarapan serta konsumsi sayur dan buah. *Action: Aceh Nutrition Journal*, 5(1), 55–63. <https://doi.org/10.30867/action.v5i1.203>
- Beal, T., Tumilowicz, A., Sutrisna, A., Izwardy, D., & Neufeld, L. M. (2018). A review of child stunting determinants in Indonesia. *Maternal & Child Nutrition*, 14(4), e12617. <https://doi.org/10.1111/mcn.12617>
- Damayanti, A. P., & Rahma, A. (2025). Pengaruh permainan edukatif terhadap peningkatan kesadaran gizi seimbang pada remaja. *Jurnal Gizi dan Kesehatan*, 5(1), 54–61. <https://doi.org/10.36086/jgk.v5i1.2893>
- Damayanti, E. R. (2022). Hubungan pola makan, aktivitas fisik, dan citra tubuh dengan status gizi lebih pada remaja putri di SMA Negeri 3 Tambun Selatan. *Indonesian Journal of Health Development*, 4(1), 35–45.
- Farikhah, N. (2021). Pengaruh media edukasi gizi berbasis infografis dan web terhadap pengetahuan dan perilaku makan gizi seimbang mahasiswa IAIN Kudus. Dalam *Proceedings of the National Conference of Islamic Natural Science (NCOINS)* (Vol. 1, No. 1, hlm. 154–168).
- Handayani, L., Saputri, M. E., Trisnowati, H., Novianti, T. N., Fitriani, I., Suryana, A. A., Vergawita, T., Nasir, A. J., & Aisyahrani, A. I. B. (2024). Flip chart-based nutrition education to improve knowledge and attitude toward stunting among adolescent girls. *Media Publikasi Promosi Kesehatan Indonesia*, 7(12), 2968–2975. <https://doi.org/10.56338/mppki.v7i12.6381>
- Nurchayani, I. D. (2020). Intervensi penyuluhan gizi seimbang dengan media video terhadap perubahan asupan zat gizi remaja putri. *Jurnal Ilmiah Kesehatan (JIKA)*, 2(3), 136–143.
- Putri, T. P., Sitorus, R. J., & Rahmiwati, A. (2023). Hubungan pengetahuan tentang gizi seimbang dengan status gizi pada remaja. *Jurnal Ners*, 7(2), 952–957. <https://doi.org/10.31004/jn.v7i2.15646>
- Romadhon, W. A., Wulandari, N. A., & Mahmudah, N. A. (2025). Peer group nutrition education for adolescent girls as a strategy to prevent stunting. *Journal of Community Service for Health*, 6(2), 41–47. <https://doi.org/10.26699/jcsh.v6i2.1261>
- Wahyuni, F. C., Karomah, U., Basrowi, R. W., Sitorus, N. L., & Lestari, L. A. (2023). Literasi gizi dan pencegahan stunting: Sebuah tinjauan sistematis. *Amerta Nutrition*, 7(3), 71–85. <https://doi.org/10.20473/amnt.v7i3.2023.71-85>
- Akombi-Inyang, B. (2025). Nutrition Programs and Policies for Maternal and Child Health. *Nutrients*, 17(15). Scopus. <https://doi.org/10.3390/nu17152397>
- Anyati, C., Okello, D. M., Mainimo, E. N., & Okello-Uma, I. (2025). Nutritional knowledge, attitude, and practices among caregivers and nutritional status of children 6–24 months: Evidence from Amuria and Soroti districts of Uganda. *Discover Food*, 5(1). Scopus. <https://doi.org/10.1007/s44187-025-00369-8>
- Azhar, M., Yasin, R., Hanif, S., Bughio, S. A., Das, J. K., & Bhutta, Z. A. (2025). Nutritional Management of Low Birth Weight and Preterm Infants in Low- and Low Middle-Income Countries. *Neonatology*, 122, 209–223. Scopus. <https://doi.org/10.1159/000542154>

- Bezie, M. M., Seifu, B. L., Asebe, H. A., Asnake, A. A., Negussie, Y. M., Asmare, Z. A., Melkam, M., & Fente, B. M. (2025). Population attributable fraction of risk factors for anemia among children aged 6-59 months: A multilevel complex data analysis using modified poisson regression model in Sub-Saharan Africa. *PLOS ONE*, 20(6 June). Scopus. <https://doi.org/10.1371/journal.pone.0324835>
- Brown, C., Kandpal, E., Lee, J., & Williams, A. (2025). Nutritional Inequality and Policy Targeting in South Asia. *Population and Development Review*. Scopus. <https://doi.org/10.1111/padr.70035>
- Burger, R., Du Plessis, L., Gangaidzo, T., Slemming, W., & Wills, G. (2025). Priority interventions for reducing stunting in the complementary feeding age group in South Africa. *Development Southern Africa*. Scopus. <https://doi.org/10.1080/0376835X.2025.2596786>
- Bustos, M., Lau, L., Manguerra, H., & Dodd, W. (2025). Sociodemographic factors associated with concurrent stunting and wasting among children experiencing extreme poverty in the Philippines: A cross-sectional study. *Nutrition and Health*, 31(2), 585–596. Scopus. <https://doi.org/10.1177/02601060231203422>
- Fahmida, U., Htet, M. K., Ameline, A. S., Angelin, T. C., Kolopaking, R., Davies-Kershaw, H., & Ferguson, E. (2025). Nutrition and non-nutrition determinants of maternal mental disorders, stress, and depression in pregnancy amidst a global pandemic: Findings from Action Against Stunting Hub Indonesia. *British Journal of Nutrition*. Scopus. <https://doi.org/10.1017/S0007114525105515>
- Fatkuriyah, L., Nafista, U. F., Hidayati, A., & Cahyono, H. D. (2025). Risk Factors of Stunting among Children in Rural Area of Jember, Indonesia. *Malaysian Journal of Nursing*, 16(4), 165–176. Scopus. <https://doi.org/10.31674/mjn.2025.v16i04.016>
- Fogarty, B., García-Martínez, A., Chawla, N. V., & Serván-Mori, E. (2025). Social and economic predictors of under-five stunting in Mexico: A comprehensive approach through the XGB model. *Journal of Global Health*, 15. Scopus. <https://doi.org/10.7189/JOGH.15.04065>
- Guo, W., Zgambo, M., Sanmei, S., & Shimpuku, Y. (2025). Prevalence and trends of coexisting forms of malnutrition and its associated factors among children aged 6–59 months in South and Southeast Asia, 1996–2022: A cross-sectional time series study. *BMC Public Health*, 25(1). Scopus. <https://doi.org/10.1186/s12889-025-23482-w>
- Haniarti, H., Sabriani, S., Nurlinda, N., & Umar, F. (2025). Socio-Cultural Influences on Stunting Children Aged 24-59 Months: A Cross-Sectional Study in the Bugis Community of Indonesia. *Media Publikasi Promosi Kesehatan Indonesia*, 8(8), 818–830. Scopus. <https://doi.org/10.56338/mppki.v8i8.7946>
- Hasib Joarder, M. A., Saha, P., Chakraborty, S., Akter, K., Amir, S., Chowdhury, M. R., Dowllah, I. M., Alam, U. K., Jai Maug, A. K., & Alam, M. M. (2025). Nutritional status of adolescents undergoing tuberculosis treatment in urban Bangladesh: Prevalence and determinants of malnutrition. *International Journal of Adolescent Medicine and Health*, 37(5), 315–323. Scopus. <https://doi.org/10.1515/ijamh-2025-0064>
- Hu, M., Liu, L., Zhang, Y., Zeng, T., Zhou, Z., Nong, Z., Qin, D.-H., Guan, H.-B., & Zeng, D.-Y. (2025). Recent trends and influencing factors of stunting among infants and young children aged 6-23 months in rural areas of Liuzhou from 2017 to 2023. *Modern Preventive Medicine*, 52(15), 2769–2774. Scopus. <https://doi.org/10.20043/j.cnki.MPM.202501052>
- Jaleel, A., Surya Goud, C. S., Shankar, S., & Venkatesh, K. (2025). Nourishing the future: Exploring the factors influencing minimum diet diversity and minimum acceptable diet among Indian children aged 6–23 months. *Journal of Public Health (Germany)*, 33(5), 1057–1069. Scopus. <https://doi.org/10.1007/s10389-023-02085-y>
- Lundberg, K., Moragues-Faus, A., Thornton, L., Cohen, N., Diekmann, L., & De-Regil, L. M. (2025). Paving the way: Urban Health, Food Systems, and the Imperative for Holistic

- City-Led Action. *F1000Research*, 14. Scopus. <https://doi.org/10.12688/f1000research.163042.1>
- Mekuriaw, B. Y., Molla, A., Negash, M., Teferi, T., Habtamu, E., Wale, M., & Chekol, A. (2025). Predictors of depression among caregivers of children with malnutrition in Gedeo zone public hospitals, Southern Ethiopia: Case-control study. *BMC Psychiatry*, 25(1). Scopus. <https://doi.org/10.1186/s12888-025-06959-1>
- Mokori, A., Kirimi, N., Ndungutse, A. H., Fusheini, Z., & Ssenyondo, M. (2025). Simultaneous Wasting and Stunting (WaSt), Wasting and Anemia (WaAn) and Wasting, Stunting and Anemia (WaStAn) Among Children 6–59 Months in Karamoja, Uganda. *Food Science and Nutrition*, 13(11). Scopus. <https://doi.org/10.1002/fsn3.71149>
- Mukhopadhyay, K. (2025). Nourishing the Future Generations: Role of Maternal Education, Empowerment and Autonomy. *Indian Pediatrics*, 62(7), 547–548. Scopus. <https://doi.org/10.1007/s13312-025-00095-5>
- Pasha, M., Shimul, M. M. H., Hossain, R., Khandker, S., Khan, S., & Mehriban, N. (2025). Nutritional status and determinants of malnutrition among under-five children in a specialized hospital in Bangladesh during the COVID-19 pandemic. *Clinical Nutrition Open Science*, 64, 192–202. Scopus. <https://doi.org/10.1016/j.nutos.2025.10.008>
- Samuel, A., Noor, R., Tefera, N., Hussien, A., Gebremedhin, G., Mbuya, M., Amare, E., Woldeyohannes, M., Fikresilassie, G., Girma, M., Petros, A., Senbeta, A., Tessema, B., Challa, F., Baye, K., Laillou, A., Chuko, T., Tollera, G., Hailu, M., ... Tessema, M. (2025). Potential Impact of Large-Scale Food Fortification in Ethiopia: Coverage and Knowledge of Fortifiable Foods. *Maternal and Child Nutrition*. Scopus. <https://doi.org/10.1111/mcn.70055>
- Shiyami, N. D. S., & Mardiana, n. (2025). RISK FACTOR FOR STUNTING IN TODDLERS AGED 24–59 MONTHS IN KERSANA PUBLIC HEALTH CENTER WORKING AREA. *Media Gizi Indonesia*, 20(1), 63–73. Scopus. <https://doi.org/10.20473/mgi.v20i1.63-73>
- Soharwardi, M. A., Malik, N. I., Anjum, R., Haleem, M. S., Leghari, I. U., Ahmad, J. B., Maryam, R., Nazir, M., Fatima, S., Ahmed, F., & Tang, K. (2025). Role of Maternal Empowerment in Addressing Child Malnutrition: Evidence from Asian Developing Countries. *Children*, 12(5). Scopus. <https://doi.org/10.3390/children12050597>
- Worku, M. G., Mohanty, I., Mengesha, Z., & Niyonsenga, T. (2025). Risk Factors of Standalone and Coexisting Forms of Undernutrition Among Children in Sub-Saharan Africa: A Study Using Data from 26 Country-Based Demographic and Health Surveys. *Nutrients*, 17(2). Scopus. <https://doi.org/10.3390/nu17020252>
- Zemariam, A. B., Abate, B. B., Alamaw, A. W., Lake, E. S., Yilak, G., Ayele, M., Tilahun, B. D., & Ngusie, H. S. (2025). Prediction of stunting and its socioeconomic determinants among adolescent girls in Ethiopia using machine learning algorithms. *PLOS ONE*, 20(1 January). Scopus. <https://doi.org/10.1371/journal.pone.0316452>

Copyright Holder :

© Siska Indrayani et.al (2025).

First Publication Right :

© Journal of World Future Medicine, Health and Nursing

This article is under: