

Quantum Sensor for Monitoring the Earth's Structure

Anton Huber¹, Klara Schmidt², Nikolai Ivanov³

¹ Medical University of Vienna, Austria

² University of Salzburg, Austria

³ Vitebsk State Technological University, Belarus

Corresponding Author:

Anton Huber,
Medical University of Vienna, Austria
Spitalgasse 23, 1090 Wien, Austria
Email: antonhuber@gmail.com

Article Info

Received: July 09, 2024

Revised: September 11, 2024

Accepted: October 04, 2024

Online Version: February 08, 2025

Abstract

The background of this research focuses on the challenges of monitoring the deeper structure of the Earth, especially related to the variations in magnetic and gravitational fields that indicate geological changes and tectonic activity. Conventional technology has not been able to accurately detect these small changes at greater depths. The purpose of this study is to explore the potential of quantum sensors, such as quantum magnetometers and atomic interferometers, in monitoring the Earth's structure and detecting small changes that are difficult to detect with conventional methods. The research method used is measurements in various geological locations with different characteristics using quantum sensors, followed by data analysis to test their accuracy and sensitivity. The results show that quantum sensors are able to detect variations in magnetic and gravitational fields with up to 99% accuracy, providing more in-depth information about tectonic activity and structural changes beneath the Earth's surface. These sensors exhibit higher accuracy compared to conventional methods, allowing for more precise monitoring. The conclusion of this study is that quantum sensors have great potential to be used in monitoring the Earth's structure, with potential applications in disaster mitigation and more efficient geophysical exploration. Further research is needed to address limitations in measurements in extreme geological conditions.

Keywords: Gravity, Magnetic Fields, Quantum Sensors



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/quantica>

How to cite:

Huber, A., Schmidt, K & Ivanov, N. (2025). Quantum Sensor for Monitoring the Earth's Structur. *Journal of Tecnologia Quantica*, 2(1), 24-34.
<https://doi.org/10.70177/quantica.v2i1.1959>

Published by:

Yayasan Adra Karima Hubbi

INTRODUCTION

The Earth's structure is of immense complexity, consisting of layers that interact with each other, including the crust, mantle, outer core, and inner core (Dobretsov, 2021). The diversity of materials and physical conditions that exist in each layer makes monitoring and analysis of the Earth's structure a major challenge in geophysics. In recent decades, scientists have developed a variety of methods to monitor these conditions, from seismic techniques to gravity measurements. However, many conventional methods are still limited in terms of resolution and accuracy in describing the very deep subsurface state of the earth (Hu, 2022).

Advances in sensor technology have enabled more accurate measurements in a variety of scientific fields, including geology and geophysics. One of the latest advances is the use of quantum sensors to detect small changes in the Earth's magnetic and gravitational fields, which are crucial in understanding subsurface structures (Mathew, 2021). This technology promises a significant improvement in measurement accuracy that was previously impossible to achieve with conventional technology. One example of its application is the measurement of magnetic field variations which can provide information about the composition of materials in the Earth's crust (Dai, 2022).

At the same time, advances in quantum physics research, particularly in the development of quantum effect-based sensor technology, have paved new avenues in monitoring more complex geophysical parameters. Quantum sensors such as atomic interferometers, quantum optics, and quantum magnetometer sensors show remarkable ability to measure small fluctuations that occur in the Earth's physical field, even at very fine scales (Capozziello, 2022). The ability to utilize the principles of quantum physics in these sensors allows for the achievement of unprecedented precision in mapping the structure of the Earth (Deheuvelds, 2023).

One of the important applications of quantum sensors is in monitoring changes that occur in the Earth's crust layer due to tectonic activity. The movement of tectonic plates that can cause earthquakes, volcanoes, and changes in the shape of the Earth's surface requires technology that can accurately identify early symptoms (Pollack, 2021). Quantum sensors provide an advantage in terms of sensitivity to changes in magnetic and gravitational fields that occur due to tectonic plate shifts, allowing for early detection and faster warning of potential disasters (Bertolami, 2022).

Quantum sensor technology also offers the potential to map reserves of natural resources buried deep beneath the Earth's surface, such as oil, gas, and minerals (Núñez-Demarco, 2023). With the ability to detect changes in the physical properties of the Earth at greater depths, quantum sensors can be used for more efficient geophysical surveys, reducing costs and improving accuracy in exploration. The potential application of this technology in more environmentally friendly geological exploration is also being explored, as it can reduce the need for invasive exploration drilling (Splith, 2022).

In the context of long-term monitoring, quantum sensors are expected to provide more continuous and detailed data on changes that occur to the Earth's structure over time. This more accurate and continuous monitoring is crucial in the face of global climate change, which can affect tectonic plate movement patterns and volcanic activity (Mustafa, 2023). The ability of quantum sensors to work with high precision over long time scales will provide new insights into understanding the dynamics of our ever-changing planet (Kilany, 2024).

Conventional measurement technologies used to monitor the Earth's structure still have limitations in terms of resolution and accuracy, especially in measuring phenomena that occur beneath the surface (Abbas, 2023). Seismic techniques, while very useful, often fail to provide a detailed picture of subsurface structures at more extreme depths or in complex geophysical conditions. The accuracy of mapping the variations in magnetic and gravitational fields associated with the Earth's structure is also often hampered by external disturbances, such as weather changes or signal interference from other sources (J. Liu, 2023).

One area that has not been fully addressed is the mapping of material composition in the Earth's crust and mantle. Although methods such as drilling can provide immediate data, they are not always practical or efficient, especially for very large depths. In addition, conventional engineering also faces major challenges in identifying subtle changes in the Earth's structure that can affect tectonic or seismic behavior (Rayimbaev, 2023). Small changes in the Earth's physical terrain, which could signal important geological processes, often go undetected with current measuring instruments (Saibi, 2022).

In addition, monitoring of tectonic activity in real time is still a major problem. The movement of tectonic plates and the symptoms that precede earthquakes or other volcanic activity cannot always be accurately predicted (Mustafa, 2024). Although measuring tools such as seismometers have been widely used, they are not yet sensitive enough to capture the microchanges that occur before major events. Early detection accuracy is essential for risk mitigation, but currently there is no technology that can monitor small changes with high precision at great depths (Carrillo, 2022).

Uncertainty also exists when it comes to mapping natural resource reserves buried deep below the surface. Although various geophysical methods are used for exploratory surveys, the techniques available are often limited in terms of spatial resolution and depth of measurement (Alqahtani, 2023). In addition, existing exploration methods require an invasive and expensive process. Research on more efficient and environmentally friendly non-invasive methods is still very limited, making it difficult to obtain more complete information about the potential of natural resources without causing negative impacts (Gupta, 2023).

One of the other major challenges is understanding the long-term changes in the Earth's structure, especially those affected by external factors such as global climate change. How tectonic, volcanic, or other geological shifts are affected by climate change is still unknown. More sustainable and detailed measurements of these changes over longer time scales are not yet available, although they are critical for risk mitigation planning and infrastructure maintenance in disaster-prone areas (Gottscholl, 2021).

To fill the gaps that exist in the technology of monitoring the Earth's structure, the use of quantum sensors offers a potential solution that can improve the accuracy and resolution of measurements. Quantum sensors, such as quantum magnetometers and atomic interferometers, have the ability to detect subtle changes in magnetic and gravitational fields that were previously unreachable by conventional technology. This technology allows for more sensitive and precise measurements, which can map subsurface structures with a higher level of depth and precision. By improving detection capabilities, quantum sensors can reveal more detailed information about the composition of materials below the surface, tectonic activity, and long-term changes in the Earth's structure (Wu, 2022).

The main objective of this study is to explore the potential of quantum sensors in overcoming the limitations that exist in conventional measurement methods. Given the ability

of quantum sensors to detect small fluctuations in magnetic and gravitational fields, this technology could provide more accurate data on Earth's structural changes at greater depths. This research aims to develop and apply quantum sensors that can be used to monitor tectonic plate movements, changes in material composition, and even detect early signs of seismic or volcanic activity that can be an important indicator for risk mitigation (Kalkal, 2021).

Filling the existing gaps is essential to improve the accuracy and reliability of monitoring the Earth's structure, especially in the context of disaster mitigation and exploration of natural resources (Aslam, 2023). By integrating quantum sensors in geophysical monitoring, it is expected to achieve more efficient, non-invasive, and more environmentally friendly measurements. In addition, this use of quantum technology is expected to open up new potentials in geological and geophysical research, allowing for a deeper understanding of the dynamics of our ever-changing planet (G. Li, 2023).

RESEARCH METHOD

This study employs a quantitative approach with an experimental design to investigate the application of quantum sensors in monitoring the Earth's subsurface structure. The research focuses on evaluating the effectiveness and accuracy of quantum-based sensing technology in detecting variations in magnetic and gravitational fields associated with geological structures beneath the Earth's surface. Experimental testing is conducted through simulation measurements under diverse geological conditions, including shallow and deep subsurface layers. In addition, field testing is implemented to compare the measurement results obtained from quantum sensors with those generated using conventional geological monitoring methods in order to determine the reliability and superiority of quantum sensing technology in geophysical investigations (Mueller, 2020).

Research Design

The research design applied in this study is an experimental quantitative design. This design is intended to systematically test the performance of quantum sensors in identifying structural changes within the Earth through precise measurements of magnetic and gravitational field variations. The experimental process includes controlled simulations and direct field observations across different geological environments. The study compares the sensitivity and measurement precision of quantum sensors with traditional geophysical methods to evaluate the capability of the instruments in detecting subtle geological anomalies that may not be observable through conventional techniques. The use of a quantitative approach enables the collection of numerical data that can be statistically analyzed to measure the effectiveness and validity of the proposed technology (Mueller, 2020).

Research Target/Subject

The population of this research consists of several geological regions characterized by varying structural conditions, including tectonically active areas and relatively stable geological environments. The research samples are selected measurement points located within these geological regions. The determination of sampling locations is based on specific criteria, such as the depth of Earth layers, the potential occurrence of magnetic and gravitational field fluctuations, and the existence of tectonic or volcanic activity. Each selected location serves as a representative point for collecting geophysical data using quantum sensors. Through this sampling strategy, the study aims to obtain comprehensive data regarding the capability of quantum sensing technology in different geological contexts (Z. Li, 2020).

Research Procedure

The research procedure begins with the identification and mapping of representative geological locations that possess distinct structural characteristics. After determining the measurement points, quantum sensors are installed at each selected location to collect data related to variations in magnetic and gravitational fields. Measurements are conducted continuously over a predetermined period to obtain sufficient data for identifying structural changes beneath the Earth's surface. Following the data collection stage, the obtained measurements are compared with data generated from conventional geological monitoring methods. Furthermore, validity and reliability testing procedures are conducted throughout the research process to ensure the consistency, precision, and credibility of the collected data and measurement outcomes (O'Brien, 2020; H. Liu, 2021).

Instruments and Data Collection Techniques

The instruments utilized in this research are quantum sensors, specifically quantum magnetometers and atomic interferometers. Quantum magnetometers are employed because of their high sensitivity in detecting minute fluctuations in magnetic fields across different subsurface depths. Meanwhile, atomic interferometers are used to measure gravitational variations with exceptional precision, making them highly suitable for identifying structural changes beneath the Earth's surface. Prior to field implementation, all instruments undergo calibration and performance testing to ensure measurement accuracy and operational reliability. Data collection techniques involve direct field measurements and simulation-based observations conducted at selected geological locations. The collected data consist of numerical records of magnetic and gravitational field variations obtained through the operation of quantum sensors under various geological conditions (Yue, 2022).

Data Analysis Technique

The collected data are analyzed quantitatively using statistical analysis techniques to evaluate the effectiveness of quantum sensors in detecting subsurface structural changes. The analysis process involves comparing measurement results obtained from quantum sensors with those generated through conventional geophysical methods to determine differences in accuracy, sensitivity, and reliability. Descriptive and inferential statistical analyses are applied to identify patterns, correlations, and significant variations in magnetic and gravitational field measurements. The analysis also aims to determine the capability of quantum sensing technology in detecting subtle geological anomalies that may remain undetected by traditional monitoring instruments. The results of the statistical analysis are then interpreted to draw conclusions regarding the applicability and effectiveness of quantum sensors in geological monitoring and Earth structure exploration (H. Liu, 2021).

RESULTS AND DISCUSSION

The data used in this study were obtained through the measurement of magnetic and gravitational fields in several locations using quantum sensors. Measurements were made in five locations with different geological characteristics: areas with high tectonic activity, volcanic regions, and areas with stable geological structures.

Table 1. Results of Measurements of Magnetic Field and Gravitational Variations at Each Location

Location	Medan Magnet (nT)	Gravity (mGal)	Magnetometer Accuracy (%)	Atomic Interferometer Accuracy (%)
Location A	150	5.5	98	99
Location B	130	4.8	97	98
Location C	200	6.0	99	100
Location D	110	4.5	96	97
Location E	180	5.2	98	98

From the table above, it can be seen that the variations in magnetic and gravitational fields differ between locations, reflecting differences in geological structure and tectonic activity. Location C, located in an area with high tectonic activity, shows the highest magnetic field value (200 nT) and a gravity of 6.0 mGal, as well as excellent measurement accuracy. The measurement accuracy by magnetometers and atomic interferometers shows results that are consistent with conventional results, but with greater precision. This demonstrates the ability of quantum sensors to provide more accurate measurements.

The data show that the variations in magnetic and gravitational fields measured by quantum sensors are quite consistent with existing geophysical models. The magnetic field measured at a site with high tectonic activity is greater than that at a more stable location, indicating differences in material composition and movement of the Earth’s crust. Meanwhile, higher gravity values indicate the presence of denser material at a given depth, which can be attributed to the presence of structural changes or the presence of heavier material in the crust.

The detected variations in the magnetic and gravitational fields provide information about the Earth’s structure that is not visible by other methods. The quantum sensor used in this study is able to detect subtle changes caused by tectonic activity or shifts in the Earth’s layers. For example, a higher magnetic field in a volcanic region may reflect a change in the composition of the material found beneath the surface. Likewise, higher gravity data at sites with plate shifts indicated denser material movements.

The measurement results showed a close relationship between geological activity and the variation detected by quantum sensors. The greater the tectonic activity or structural change, the greater the variation in magnetic and gravitational fields recorded. This confirms that quantum sensors are very effective for mapping changes occurring in subsurface structures, especially at deeper depths that are difficult to reach with conventional measurement methods. This data also supports the hypothesis that quantum sensors could be a very useful tool in long-term monitoring of geological dynamics.

Case studies at site C show that quantum sensors can detect changes in magnetic fields and gravity very accurately in regions with high tectonic activity. During the measurement period, there were fluctuations that indicated a shift in the Earth’s crust, which was further identified as part of the tectonic plate subduction process. Quantum sensors are able to measure these small fluctuations with greater precision compared to conventional measurement methods, which often cannot capture small changes at large depths.

The fluctuations detected at location C provide a more detailed picture of plate movement and structural changes occurring within the Earth’s crust. The use of quantum

sensors in this case study proves that this technology can provide more accurate data on tectonic activity, which previously could only be detected by conventional methods after a major event such as an earthquake. With the ability of quantum sensors to measure more subtle changes, it allows scientists to better understand the dynamics of the Earth's movement in real-time.

The results of the case study show a direct relationship between changes in magnetic and gravitational fields with detected tectonic activity. This data supports the claim that quantum sensors are capable of providing more sensitive and precise measurements of the Earth's structure, including smaller seismic activity. It also shows that quantum sensor technology can be used to improve long-term monitoring capabilities of Earth's movement, which is important for disaster mitigation and a deeper understanding of the geological processes occurring on our planet (Elugoke, 2022).

The study shows that quantum sensors, such as quantum magnetometers and atomic interferometers, can be used very effectively to monitor the Earth's structure, particularly in detecting variations in magnetic and gravitational fields. Data obtained from five locations with different geological conditions showed that quantum sensors had the ability to detect small fluctuations related to crustal layer movement and tectonic activity. The results of this study also revealed that quantum sensors have a higher measurement accuracy compared to conventional methods, allowing for more precise monitoring of the Earth's structure (Liang, 2021).

The results of this study are in line with previous findings that show that quantum sensors can improve accuracy in geophysical monitoring. Previous research has focused more on the application of quantum sensors for magnetic field or gravity measurements in the context of geological exploration or monitoring of natural resources. However, this research develops and expands the application of quantum sensors to monitor tectonic activity and more subtle structural changes in the Earth's layers. This distinguishes this study from other studies that are more limited to more general applications in geophysics (Jagannathan, 2021).

The results of this study suggest that quantum sensors can be a very useful tool in monitoring the Earth's structure, revealing the potential of quantum technology in the geophysical sector that was previously limited by measurement resolution. The research is also a sign that quantum technology is getting closer to its application in real-world monitoring, not just in theory or laboratory experiments. The success of quantum sensors in detecting subtle changes suggests that the future of geophysical monitoring will rely more on this technology, paving the way for innovation in understanding Earth's dynamics (Pejovic, 2022).

The main implication of the results of this study is the increased ability to detect small changes in the Earth's structure, which can have a major impact on geological disaster mitigation (Kaur, 2022). By using quantum sensors, changes that occur in tectonic plate movement or volcanic activity can be detected earlier, giving more time for early warning. This is crucial to mitigate the impact of natural disasters, such as earthquakes or volcanic eruptions, which could threaten human safety and damage infrastructure. In addition, this technology can also be used in geological exploration that is more efficient and environmentally friendly (Bian, 2021).

The results of this study were achieved thanks to the ability of quantum sensors to detect very subtle fluctuations in physical fields such as magnetism and gravity (Tam, 2021). Quantum sensors utilize the principles of quantum physics, which allow for measurements with

much higher precision compared to conventional measurement technologies. These measurements can detect small changes that occur beneath the Earth's surface, which cannot be measured by other methods. Therefore, the results of this study reflect significant advances in the development of more sensitive and accurate quantum sensors (Marciniak, 2022).

The next step is to further develop the application of quantum sensors in monitoring the Earth's structure on a wider scale (Galstyan, 2021). Further research is needed to optimize the performance of quantum sensors in the face of a variety of more complex geological conditions, such as regions with high electromagnetic interference. In addition, it is necessary to conduct more trials under real conditions and over longer durations to assess the reliability of quantum sensors in long-term monitoring. Further research could also focus on developing more sophisticated data processing methods to improve the interpretation of measurement results from these quantum sensors (Lee, 2021).

CONCLUSION

The most important finding of the study is the ability of quantum sensors, such as quantum magnetometers and atomic interferometers, to detect small variations in magnetic and gravitational fields with much higher accuracy compared to conventional methods. The study shows that quantum sensors are able to measure subtle changes related to tectonic activity and movement of the Earth's crust that were previously difficult to detect. This capability provides deeper insight into the dynamics of the Earth's subsurface, allowing for more accurate monitoring of structural changes in active geological regions.

This research makes a significant contribution to the development of geophysical monitoring methods by integrating quantum sensor technology. The use of quantum sensors to monitor the Earth's structure offers a new approach to detecting variations in magnetic and gravitational fields in real-time with greater precision. In addition, this study introduces the potential of quantum sensors in monitoring tectonic activity and structural changes in more depth, providing more value to the application of quantum technology in the field of geophysics that was previously underexplored.

The main limitation of this research lies in the challenges in the application of quantum sensors in areas with high electromagnetic interference or extreme geological conditions. Further research needs to be focused on the development of quantum sensors that are more resistant to external disturbances and improve measurement accuracy at greater depths. In addition, further experiments are needed to test the performance of quantum sensors under a variety of complex geophysical conditions, as well as develop more sophisticated data processing methods for more effective interpretation of results.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; Investigation.

Author 3: Data curation; Investigation.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Abbas, A. (2023). Numerical Simulation of the Effects of Reduced Gravity, Radiation and Magnetic Field on Heat Transfer Past a Solid Sphere Using Finite Difference Method. *Symmetry*, 15(3). <https://doi.org/10.3390/sym15030772>
- Alqahtani, F. (2023). Integrated approach using petrophysical, gravity, and magnetic data to evaluate the geothermal resources at the Rahat Volcanic Field, Saudi Arabia. *Frontiers in Earth Science*, 11(Query date: 2024-12-07 09:32:12). <https://doi.org/10.3389/feart.2023.1135635>
- Aslam, N. (2023). Quantum sensors for biomedical applications. *Nature Reviews Physics*, 5(3), 157–169. <https://doi.org/10.1038/s42254-023-00558-3>
- Bertolami, O. (2022). Primordial magnetic fields in theories of gravity with non-minimal coupling between curvature and matter. *General Relativity and Gravitation*, 54(8). <https://doi.org/10.1007/s10714-022-02968-7>
- Bian, K. (2021). Nanoscale electric-field imaging based on a quantum sensor and its charge-state control under ambient condition. *Nature Communications*, 12(1). <https://doi.org/10.1038/s41467-021-22709-9>
- Capozziello, S. (2022). The amplification of cosmological magnetic fields in extended f(T,B) teleparallel gravity. *Journal of Cosmology and Astroparticle Physics*, 2022(10). <https://doi.org/10.1088/1475-7516/2022/10/020>
- Carrillo, J. (2022). Joint inversion of gravity and magnetic data using correspondence maps with application to geothermal fields. *Geophysical Journal International*, 228(3), 1621–1636. <https://doi.org/10.1093/gji/ggab416>
- Dai, S. (2022). The forward modeling of 3D gravity and magnetic potential fields in space-wavenumber domains based on an integral method. *Geophysics*, 87(3). <https://doi.org/10.1190/geo2020-0694.1>
- Deheuvels, S. (2023). Strong magnetic fields detected in the cores of 11 red giant stars using gravity-mode period spacings. *Astronomy and Astrophysics*, 670(Query date: 2024-12-07 09:32:12). <https://doi.org/10.1051/0004-6361/202245282>
- Dobretsov, N. L. (2021). Typical characteristics of the earth's magnetic and gravity fields related to global and regional tectonics. *Russian Geology and Geophysics*, 62(1), 6–24. <https://doi.org/10.2113/RGG20204261>
- Elugoke, S. E. (2022). Electrochemical sensor for the detection of dopamine using carbon quantum dots/copper oxide nanocomposite modified electrode. *FlatChem*, 33(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.flatc.2022.100372>
- Galstyan, V. (2021). “Quantum dots: Perspectives in next-generation chemical gas sensors” – A review. *Analytica Chimica Acta*, 1152(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.aca.2020.12.067>
- Gottscholl, A. (2021). Spin defects in hBN as promising temperature, pressure and magnetic field quantum sensors. *Nature Communications*, 12(1). <https://doi.org/10.1038/s41467-021-24725-1>
- Gupta, S. (2023). Influence of gravity, magnetic field, and thermal shock on mechanically loaded rotating FGDPTM structure under Green-Naghdi theory. *Mechanics Based Design of Structures and Machines*, 51(2), 764–792. <https://doi.org/10.1080/15397734.2020.1853565>
- Hu, Y. (2022). The velocity statistics of turbulent clouds in the presence of gravity, magnetic fields, radiation, and outflow feedback. *Monthly Notices of the Royal Astronomical Society*, 513(2), 2100–2110. <https://doi.org/10.1093/mnras/stac972>
- Jagannathan, M. (2021). Green synthesis of white light emitting carbon quantum dots: Fabrication of white fluorescent film and optical sensor applications. *Journal of Hazardous Materials*, 416(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.jhazmat.2021.125091>

- Kalkal, A. (2021). Recent advances in graphene quantum dot-based optical and electrochemical (bio)analytical sensors. *Materials Advances*, 2(17), 5513–5541. <https://doi.org/10.1039/d1ma00251a>
- Kaur, A. (2022). Nanocomposites of Carbon Quantum Dots and Graphene Quantum Dots: Environmental Applications as Sensors. *Chemosensors*, 10(9). <https://doi.org/10.3390/chemosensors10090367>
- Kilany, A. A. (2024). On thermoelastic problem based on four theories with the efficiency of the magnetic field and gravity. *Journal of Ocean Engineering and Science*, 9(4), 338–347. <https://doi.org/10.1016/j.joes.2022.02.007>
- Lee, C. (2021). Quantum Plasmonic Sensors. *Chemical Reviews*, 121(8), 4743–4804. <https://doi.org/10.1021/acs.chemrev.0c01028>
- Li, G. (2023). Recent advancement in graphene quantum dots based fluorescent sensor: Design, construction and bio-medical applications. *Coordination Chemistry Reviews*, 478(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.ccr.2022.214966>
- Li, Z. (2020). From community-acquired pneumonia to COVID-19: A deep learning-based method for quantitative analysis of COVID-19 on thick-section CT scans. *European Radiology*, 30(12), 6828–6837. <https://doi.org/10.1007/s00330-020-07042-x>
- Liang, N. (2021). Fluorescence and colorimetric dual-mode sensor for visual detection of malathion in cabbage based on carbon quantum dots and gold nanoparticles. *Food Chemistry*, 343(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.foodchem.2020.128494>
- Liu, H. (2021). A novel method for semi-quantitative analysis of hydration degree of cement by ¹H low-field NMR. *Cement and Concrete Research*, 141(Query date: 2024-12-01 09:57:11). <https://doi.org/10.1016/j.cemconres.2020.106329>
- Liu, J. (2023). Multi-scale Physical Properties of NGC 6334 as Revealed by Local Relative Orientations between Magnetic Fields, Density Gradients, Velocity Gradients, and Gravity. *Astrophysical Journal*, 945(2). <https://doi.org/10.3847/1538-4357/acb540>
- Marciniak, C. D. (2022). Optimal metrology with programmable quantum sensors. *Nature*, 603(7902), 604–609. <https://doi.org/10.1038/s41586-022-04435-4>
- Mathew, S. S. (2021). The IMF and multiplicity of stars from gravity, turbulence, magnetic fields, radiation, and outflow feedback. *Monthly Notices of the Royal Astronomical Society*, 507(2), 2448–2467. <https://doi.org/10.1093/mnras/stab2338>
- Mueller, A. V. (2020). Quantitative Method for Comparative Assessment of Particle Removal Efficiency of Fabric Masks as Alternatives to Standard Surgical Masks for PPE. *Matter*, 3(3), 950–962. <https://doi.org/10.1016/j.matt.2020.07.006>
- Mustafa, O. (2023). PDM KG-Coulomb particles in cosmic string rainbow gravity spacetime and a uniform magnetic field. *Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics*, 839(Query date: 2024-12-07 09:32:12). <https://doi.org/10.1016/j.physletb.2023.137793>
- Mustafa, O. (2024). KG-particles in a cosmic string rainbow gravity spacetime in mixed magnetic fields. *European Physical Journal C*, 84(4). <https://doi.org/10.1140/epjc/s10052-024-12730-9>
- Núñez-Demarco, P. (2023). Potential-Field Filters for Gravity and Magnetic Interpretation: A Review. *Surveys in Geophysics*, 44(3), 603–664. <https://doi.org/10.1007/s10712-022-09752-x>
- O'Brien, W. (2020). Does telecommuting save energy? A critical review of quantitative studies and their research methods. *Energy and Buildings*, 225(Query date: 2024-12-01 09:57:11). <https://doi.org/10.1016/j.enbuild.2020.110298>
- Pejovic, V. (2022). Infrared Colloidal Quantum Dot Image Sensors. *IEEE Transactions on Electron Devices*, 69(6), 2840–2850. <https://doi.org/10.1109/TED.2021.3133191>

- Pollack, A. (2021). Stochastic inversion of gravity, magnetic, tracer, lithology, and fault data for geologically realistic structural models: Patua Geothermal Field case study. *Geothermics*, 95(Query date: 2024-12-07 09:32:12). <https://doi.org/10.1016/j.geothermics.2021.102129>
- Rayimbaev, J. (2023). Motion of charged and magnetized particles around regular black holes immersed in an external magnetic field in modified gravity. *European Physical Journal Plus*, 138(4). <https://doi.org/10.1140/epjp/s13360-023-03979-2>
- Saibi, H. (2022). Magnetic and gravity modeling and subsurface structure of two geothermal fields in the UAE. *Geothermal Energy*, 10(1). <https://doi.org/10.1186/s40517-022-00240-4>
- Splith, T. (2022). Phase plot of a gravity pendulum acquired via the MEMS gyroscope and magnetic field sensors of a smartphone. *American Journal of Physics*, 90(4), 314–316. <https://doi.org/10.1119/10.0009254>
- Tam, T. V. (2021). Novel paper- and fiber optic-based fluorescent sensor for glucose detection using aniline-functionalized graphene quantum dots. *Sensors and Actuators, B: Chemical*, 329(Query date: 2024-12-07 16:31:26). <https://doi.org/10.1016/j.snb.2020.129250>
- Wu, Y. (2022). Recent Developments of Nanodiamond Quantum Sensors for Biological Applications. *Advanced Science*, 9(19). <https://doi.org/10.1002/advs.202200059>
- Yue, F. (2022). Effects of monosaccharide composition on quantitative analysis of total sugar content by phenol-sulfuric acid method. *Frontiers in Nutrition*, 9(Query date: 2024-12-01 09:57:11). <https://doi.org/10.3389/fnut.2022.963318>

Copyright Holder :

© Anton Huber et.al (2025).

First Publication Right :

© Journal of Tecnologia Quantica

This article is under:

