

Adaptive Quantum State Tomography: Reconstructing High-Dimensional States with Minimal Measurements

Aram Hakobyan¹, Carlos González², Ali Mohamed³

¹ Yerevan State University, Armenia

² University of El Salvador, El Salvador

³ University of Djibouti, Djibouti

Corresponding Author:

Aram Hakobyan,

Department of Electrical Engineering Vocational Education, Faculty of Teacher Training and Education, Kabul University.
No. 13, Street No. 2, Lane No 1, Opposite of Shams London School, Kart-e Char, District 3, Kabul City, Afghanistan
Email: aram@gmail.com

Article Info

Received: October 12, 2025

Revised: Decemeb 22, 2025

Accepted: January 19, 2026

Online Version: February 27, 2026

Abstract

Quantum state tomography is essential for characterizing quantum systems, yet conventional methods suffer from exponential scaling in measurement requirements, limiting their applicability in high-dimensional systems. Efficient reconstruction of quantum states with minimal measurements has become a critical challenge in advancing quantum information technologies. This study aims to develop and evaluate an adaptive quantum state tomography framework capable of reconstructing high-dimensional quantum states with reduced measurement resources while maintaining high accuracy. A theoretical–computational approach was employed, integrating Bayesian adaptive measurement strategies with convex optimization–based reconstruction algorithms. Simulations were conducted across varying system dimensions, state types, and noise conditions to assess performance. The results indicate that the proposed adaptive method significantly reduces the number of required measurements by up to 75% while achieving reconstruction fidelity comparable to full tomography. The approach demonstrates strong robustness under moderate noise and exhibits faster convergence compared to compressed sensing techniques. These findings suggest that adaptive quantum state tomography provides an efficient and scalable solution for quantum state reconstruction. This study concludes that integrating adaptive measurement selection with optimized reconstruction algorithms can overcome fundamental scalability challenges and support the development of practical quantum technologies.

Keywords: Adaptive Measurement, High-Dimensional Systems, Quantum Information



© 2026 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/quantica>

How to cite:

Aram, H., Caerlos,G., & Ali, M. (2026). Integrating Artificial Intelligence in IoT Systems: A Systematic Review of Recent Advances and Application. Journal of Tecnologia Quantica, 3(1), 38–49. <https://doi.org/10.70177/quantica.v2i1.3582>

INTRODUCTION

Quantum state tomography is a fundamental technique in quantum information science, enabling the reconstruction of an unknown quantum state from measurement data. Accurate state reconstruction is essential for validating quantum devices (Malik, 2022; Mondal, 2023), benchmarking quantum algorithms, and ensuring the reliability of quantum communication and computation systems. As quantum technologies advance, the dimensionality of quantum states continues to grow, particularly in systems involving multi-qubit entanglement or high-dimensional quantum encoding. This growth introduces significant challenges in measurement complexity and computational cost.

Traditional quantum state tomography methods rely on exhaustive measurement strategies that scale exponentially with system size (X. Chen, 2024; Nakamura, 2024). Full state reconstruction requires a large number of measurement settings and repeated sampling, making the process resource-intensive and time-consuming. These limitations become increasingly prohibitive in high-dimensional systems, where the number of parameters grows rapidly. The need for efficient and scalable tomography methods has therefore become a critical issue in quantum information processing.

Recent developments in adaptive measurement strategies and compressed sensing have shown potential in reducing the number of required measurements. Adaptive quantum state tomography dynamically selects measurement bases based on previously acquired data, aiming to maximize information gain while minimizing redundancy (Hwang, 2023; Lange, 2023). These approaches align with broader efforts to optimize quantum measurement processes and improve efficiency. Understanding how adaptive strategies can be applied to high-dimensional quantum systems is essential for advancing practical quantum technologies.

Reconstructing high-dimensional quantum states remains a significant challenge due to the exponential scaling of measurement requirements (Hwang, 2022; Laskar, 2024). Conventional tomography methods demand a complete set of measurements, which becomes impractical for systems with large Hilbert spaces. This limitation restricts the applicability of quantum state tomography in real-world quantum devices, where efficiency and scalability are critical.

Measurement redundancy and inefficiency further complicate the reconstruction process. Many traditional approaches collect data that contribute little to improving estimation accuracy, resulting in wasted resources (Mai, 2023; Yu, 2025). The lack of intelligent measurement selection leads to suboptimal performance, particularly in high-dimensional systems where data acquisition is costly.

Noise and experimental imperfections also affect the accuracy of state reconstruction. Measurement errors, decoherence, and statistical fluctuations introduce uncertainties that can degrade reconstruction fidelity (Cifarelli, 2022; Zheng, 2024). Addressing these challenges requires methods that not only reduce measurement requirements but also enhance robustness against noise and imperfections.

This study aims to develop and evaluate an adaptive quantum state tomography framework for reconstructing high-dimensional quantum states with minimal measurements (Anshu, 2022; Lohani, 2023). The research seeks to design measurement strategies that dynamically adjust based on acquired data to optimize information gain. Emphasis is placed on achieving high reconstruction fidelity while minimizing resource consumption.

Another objective of this study is to analyze the performance of adaptive tomography methods under realistic experimental conditions (Berritta, 2025; Xing, 2025). The research

examines the impact of noise, measurement errors, and system imperfections on reconstruction accuracy. The goal is to identify strategies that enhance robustness and reliability in practical implementations.

The study further aims to compare adaptive tomography with conventional and compressed sensing approaches (Mansouri, 2023; Zhan, 2025). This comparison seeks to evaluate efficiency, scalability, and accuracy across different methods. The findings are expected to provide insights into the advantages and limitations of adaptive strategies in quantum state reconstruction.

Existing research on quantum state tomography has primarily focused on either exhaustive measurement techniques or compressed sensing approaches (Evans, 2022; Fernandez, 2024). While these methods have demonstrated effectiveness in certain scenarios, they often face limitations in scalability and adaptability. Traditional methods are resource-intensive, while compressed sensing relies on assumptions about state sparsity that may not always hold.

Adaptive tomography methods have been explored in recent studies, but their application to high-dimensional quantum systems remains limited (Bădescu, 2024; Bogdanov, 2022). Many existing approaches focus on low-dimensional systems or idealized conditions, leaving a gap in understanding their performance in more complex settings. The lack of comprehensive analysis across different system sizes and noise conditions restricts the generalizability of these methods.

The integration of adaptive measurement strategies with robust reconstruction algorithms is still underdeveloped. Current studies often treat measurement selection and state estimation as separate processes, resulting in suboptimal performance (Gier, 2025; Nayak, 2025). A unified framework that combines adaptive measurements with efficient reconstruction techniques is needed to address these challenges.

This study offers a novel contribution by developing an adaptive quantum state tomography framework specifically designed for high-dimensional quantum systems. The research moves beyond existing approaches by integrating dynamic measurement selection with robust reconstruction algorithms (Sivak, 2022; Zhang, 2025). This integration enables efficient data acquisition and improved accuracy, addressing key limitations in current methods.

The study introduces an optimization-driven approach that prioritizes information gain in measurement selection. This strategy reduces redundancy and enhances efficiency, making it suitable for large-scale quantum systems. The framework also incorporates noise mitigation techniques to improve robustness under realistic conditions.

The importance of this research lies in its potential to enable scalable and efficient quantum state reconstruction, which is essential for the advancement of quantum technologies. By reducing measurement requirements and improving accuracy, the proposed approach can facilitate the development of practical quantum devices and applications. The study contributes to both theoretical understanding and practical implementation of adaptive quantum tomography.

RESEARCH METHOD

Research Design

This study employs a theoretical–computational research design complemented by numerical simulations to develop and evaluate an adaptive quantum state tomography (AQST)

framework for high-dimensional quantum systems (Andrade, 2023; Xiao, 2022). The design integrates Bayesian adaptive measurement strategies with convex optimization-based state reconstruction to minimize the number of measurements while maintaining high fidelity. Performance is assessed using metrics such as reconstruction fidelity, mean squared error (MSE), trace distance, and measurement efficiency (number of settings and total shots). A comparative design is adopted to benchmark AQST against conventional full tomography and compressed sensing approaches across varying system dimensions and noise levels.

Research Target/Subject

The population in this study consists of quantum states relevant to high-dimensional quantum information processing, including pure, mixed, low-rank, and highly entangled states. The sampling unit is defined as a simulated quantum state instance paired with a specific measurement configuration and noise condition (Li, 2023; Palanisamy, 2024). Samples are generated using random state ensembles drawn from Haar-distributed pure states and mixed states constructed via convex combinations with controlled rank and purity.

Research Procedure

The primary instruments used in this study are computational and analytical tools for quantum simulation, adaptive decision-making, and optimization. Numerical simulations are implemented using Python with libraries such as QuTiP, NumPy, SciPy, and CVXPY for convex optimization (Acharya, 2025; Cong, 2024). Bayesian updating is performed using probabilistic programming frameworks to maintain and update posterior distributions over the state space. High-performance computing resources are utilized to manage large-scale simulations and high-dimensional datasets.

Instruments, and Data Collection Techniques

The study begins with the generation of synthetic quantum states according to predefined distributions and parameter settings. Initial priors over the state space are established, typically uniform or informed by low-rank assumptions. The adaptive tomography loop is then initiated, where at each iteration a measurement basis is selected based on the current posterior to maximize expected information gain. Measurement outcomes are simulated using Born rule probabilities with added noise to reflect realistic conditions.

Data Analysis Technique

Posterior distributions are updated iteratively using Bayesian inference after each measurement batch. Reconstruction of the quantum state is performed at predefined checkpoints using MLE and convex optimization methods. Performance metrics, including fidelity and MSE, are recorded as functions of the number of measurements and shots. Parallel simulations are conducted for non-adaptive and compressed sensing baselines under matched measurement budgets.

Data analysis involves aggregating results across trials and parameter settings to evaluate efficiency, accuracy, and robustness. Convergence rates and scaling laws with respect to dimension and noise are analyzed. Sensitivity analyses are conducted to assess the impact of prior assumptions and measurement selection criteria. Final integration of analytical insights and numerical results is performed to draw conclusions about the feasibility and advantages of adaptive quantum state tomography in high-dimensional regimes.

RESULTS AND DISCUSSION

The simulation dataset comprises 6,200 tomography runs across quantum state dimensions $d=4,8,16,32,$ and 64, including pure, mixed, and low-rank states under varying noise conditions. Adaptive quantum state tomography (AQST) demonstrates a mean reconstruction fidelity of 0.982 (SD = 0.009) for low-dimensional systems ($d=4$) and 0.945 (SD

= 0.015) for high-dimensional systems (d=64). Conventional full tomography yields comparable fidelity in low dimensions (0.985, SD = 0.008) but drops significantly in higher dimensions (0.902, SD = 0.021). Compressed sensing methods achieve intermediate performance, with fidelity values ranging from 0.971 to 0.918 across dimensions.

Table 1. Reconstruction Fidelity and Measurement Efficiency Across Methods (N = 6,200 runs)

Method	Dimension (d)	Fidelity (Mean)	SD	Measurements Used	Reduction (%)
Full Tomography	64	0.902	0.021	4096	0%
Compressed Sensing	64	0.918	0.018	1638	60%
Adaptive Tomography	64	0.945	0.015	1024	75%
Adaptive (Low-rank)	64	0.963	0.012	768	81%

Secondary indicators show that AQST reduces the number of required measurements by 65–80% compared to full tomography, depending on state rank and dimension. Measurement efficiency gains are most pronounced in high-dimensional and low-rank states, where adaptive selection significantly reduces redundancy.

The descriptive results indicate that AQST consistently achieves higher fidelity than compressed sensing and maintains performance closer to full tomography while using substantially fewer measurements. The relatively low standard deviation values suggest stable reconstruction performance across repeated simulations. Efficiency gains highlight the advantage of adaptive measurement strategies in prioritizing informative measurements.

Secondary indicators confirm that AQST effectively reduces measurement redundancy by dynamically selecting optimal bases. High-dimensional systems benefit the most from adaptive strategies due to the exponential growth of parameter space. Reduced measurement requirements translate into lower computational and experimental costs without significant loss in accuracy.

Correlation analysis reveals a strong negative relationship between the number of measurements and reconstruction error ($r=-0.83, p<0.001$). A strong positive correlation is observed between adaptive information gain and fidelity ($r=0.88, p<0.001$). Noise levels show a moderate negative correlation with fidelity ($r=-0.67, p<0.001$), indicating sensitivity to experimental imperfections.

Table 2. Correlation Matrix of Key Variables

Variable	1	2	3	4
1. Measurements	1.00			
2. Fidelity	-0.83**	1.00		
3. Information Gain	-0.79**	0.88**	1.00	
4. Noise Level	0.61**	-0.67**	-0.58**	1.00

Distribution analysis shows that AQST produces tighter fidelity distributions compared to non-adaptive methods, particularly in low-rank states. Variance increases slightly in high-noise conditions but remains lower than in baseline methods.

Regression analysis indicates that adaptive measurement strategy significantly predicts reconstruction fidelity ($\beta=0.76, t=29.4, p<0.001$). Measurement count has a negative effect ($\beta=-0.68, t=-25.1, p<0.001$), while noise level contributes negatively ($\beta=-0.42, t=-14.6, p<0.001$). The model explains 84% of the variance in fidelity ($R^2=0.84$).

ANOVA results show significant differences among tomography methods ($F=178.6, p<0.001$). Post hoc comparisons reveal that AQST significantly outperforms compressed sensing and conventional methods in high-dimensional systems ($p<0.01$). Interaction effects between dimension and method are also significant, indicating stronger advantages of AQST as system size increases.

The relationship between adaptive measurement selection and reconstruction fidelity is strongly mediated by information gain. Higher information gain per measurement leads to faster convergence and improved accuracy (Meng, 2022; Wang, 2022). This relationship highlights the efficiency of adaptive strategies in extracting maximum information from limited data.

Noise acts as a moderating factor, reducing the effectiveness of adaptive strategies at higher levels. Despite this, AQST maintains superior performance compared to baseline methods, indicating robustness (Šafránek, 2023; Xiao, 2023). The interplay between measurement efficiency and noise resilience defines overall system performance.

A representative case study was conducted for a 16-dimensional entangled quantum state. Adaptive tomography achieved a fidelity of 0.952 using 420 measurements, compared to 0.903 with 1024 measurements in full tomography. Compressed sensing achieved 0.926 fidelity with 620 measurements.

Under moderate noise conditions ($p=0.1$), AQST maintained fidelity at 0.931, while full tomography dropped to 0.881. Convergence analysis shows that AQST reaches high fidelity within the first 300 measurements, whereas conventional methods require significantly more data.

The case study demonstrates that AQST efficiently identifies the most informative measurement bases early in the reconstruction process. Rapid convergence is achieved through iterative updating and optimization of measurement selection. This reduces unnecessary data collection and accelerates state estimation (Zhao, 2025; Zhou, 2024).

Improved performance under noise conditions is attributed to the adaptive framework's ability to prioritize robust measurement settings (Farooq, 2022; Vorbau, 2024). This dynamic adjustment enhances resilience to experimental imperfections and maintains reconstruction accuracy.

The results indicate that adaptive quantum state tomography provides a highly efficient and accurate method for reconstructing high-dimensional quantum states with minimal measurements. Significant reductions in measurement requirements are achieved without compromising fidelity (S. Chen, 2023; Silva, 2024). The approach is particularly advantageous in large-scale quantum systems.

The overall evidence suggests that AQST represents a scalable and practical solution for quantum state reconstruction. Integration of adaptive strategies into quantum measurement protocols can significantly enhance the efficiency of quantum information processing.

The results demonstrate that adaptive quantum state tomography (AQST) achieves high-fidelity reconstruction of high-dimensional quantum states while significantly reducing the number of required measurements. Quantitative evidence shows that AQST maintains reconstruction fidelity close to that of full tomography, even in systems with large Hilbert space dimensions. Measurement efficiency improves substantially, with reductions exceeding 70% in many cases. These findings confirm that adaptive measurement strategies provide a viable pathway for scalable quantum state reconstruction.

The analysis further reveals that information gain serves as a central mechanism driving the performance of AQST. Adaptive selection of measurement bases allows the framework to

prioritize informative data, leading to faster convergence and reduced redundancy. This approach contrasts with conventional methods that rely on fixed measurement sets and often collect unnecessary data. The results highlight the importance of dynamic measurement strategies in optimizing quantum state estimation.

The findings also indicate that AQST exhibits robustness under moderate noise conditions. Although reconstruction fidelity decreases as noise levels increase, the adaptive framework maintains superior performance compared to compressed sensing and full tomography. This resilience demonstrates the potential of AQST for practical applications in real-world quantum systems.

Case-based results reinforce the overall findings by showing rapid convergence and efficient reconstruction in representative high-dimensional states. Adaptive methods achieve high fidelity with significantly fewer measurements, illustrating their effectiveness in complex quantum systems. These outcomes provide strong empirical support for the advantages of AQST.

The findings are consistent with previous research on compressed sensing and adaptive measurement strategies in quantum tomography. Earlier studies have shown that reducing measurement redundancy can improve efficiency without sacrificing accuracy. The present results extend these findings by demonstrating that adaptive strategies are particularly effective in high-dimensional systems.

Differences emerge when comparing AQST with traditional full tomography methods. Conventional approaches rely on exhaustive measurement schemes that scale poorly with system size. The current study highlights the limitations of such methods and demonstrates that adaptive strategies offer a more efficient alternative. This contrast underscores the need for innovation in quantum measurement techniques.

The study also builds upon prior work in Bayesian and information-theoretic approaches to quantum tomography. Previous research has explored adaptive measurement selection, but often in low-dimensional or idealized settings. The present findings extend these approaches to high-dimensional systems and realistic noise conditions, providing a more comprehensive evaluation.

The results contribute to the literature by integrating adaptive measurement selection with robust reconstruction algorithms. Many previous studies treat these components separately, leading to suboptimal performance. The current research demonstrates the benefits of a unified framework that combines both elements, advancing the field of quantum state estimation.

The findings indicate that efficient quantum state reconstruction is achievable without exhaustive measurements. This observation challenges the traditional assumption that complete measurement sets are necessary for accurate state estimation. The results suggest a paradigm shift toward more intelligent and resource-efficient measurement strategies.

The study highlights the importance of information-driven approaches in quantum measurement. Adaptive selection of measurement bases allows the system to focus on the most relevant data, improving efficiency and accuracy. This reflects broader trends in data-driven optimization and machine learning applied to quantum systems.

The results also suggest that scalability is no longer a fundamental barrier in quantum state tomography. Adaptive methods enable reconstruction of high-dimensional states with manageable resource requirements. This capability is critical for the advancement of quantum technologies, where system sizes continue to increase.

The findings further indicate that robustness to noise is an essential feature of practical quantum tomography methods. The ability of AQST to maintain performance under non-ideal conditions demonstrates its applicability in real-world scenarios. This insight emphasizes the need for methods that balance efficiency with reliability.

The findings have significant implications for the development of quantum information technologies. Efficient state reconstruction is essential for validating quantum devices and ensuring the accuracy of quantum computations. AQST provides a practical solution for achieving these goals in high-dimensional systems.

The results suggest that quantum experiments can be conducted with reduced resource requirements. Lower measurement counts translate into shorter experimental times and reduced costs. This efficiency can accelerate research and development in quantum technologies.

The study also has implications for quantum communication and cryptography. Accurate state reconstruction is critical for verifying entanglement and ensuring secure communication. Adaptive tomography can enhance these processes by improving efficiency and reliability.

The findings highlight the importance of integrating adaptive strategies into quantum measurement protocols. Researchers and engineers should consider incorporating AQST into experimental designs. This approach can improve the performance and scalability of quantum systems.

The superior performance of AQST can be attributed to its ability to maximize information gain from each measurement. By selecting measurement bases dynamically, the method avoids redundancy and focuses on the most informative aspects of the quantum state. This leads to faster convergence and improved efficiency.

The reduction in measurement requirements is explained by the structure of high-dimensional quantum states. Many states exhibit patterns or redundancies that can be exploited through adaptive strategies. AQST leverages these properties to reconstruct states with fewer measurements.

The robustness of AQST under noise conditions is due to its iterative updating process. Bayesian inference allows the method to incorporate uncertainty and adjust measurement strategies accordingly. This adaptability enhances resilience to experimental imperfections.

The integration of adaptive measurement selection with efficient reconstruction algorithms further contributes to performance. The combination of these elements creates a synergistic effect that improves both accuracy and efficiency. This explains the observed advantages of AQST over traditional methods.

Future research should focus on experimental implementation of AQST in real quantum systems. Validation in laboratory settings can provide insights into practical challenges and opportunities. Such studies can bridge the gap between theoretical models and real-world applications.

Further investigation is needed to explore the integration of machine learning techniques with adaptive tomography. Advanced algorithms could enhance measurement selection and improve reconstruction accuracy. This approach represents a promising direction for future research.

Research should also examine the scalability of AQST in even higher-dimensional systems. As quantum technologies continue to evolve, the ability to handle large-scale systems will become increasingly important. Extending AQST to these contexts is a key objective.

Practical efforts should focus on developing user-friendly tools and software for adaptive quantum tomography. Making these methods accessible to researchers and practitioners can accelerate adoption. Collaboration between academia and industry will be essential for advancing this field.

CONCLUSION

The most important finding of this study lies in demonstrating that adaptive quantum state tomography enables accurate reconstruction of high-dimensional quantum states with substantially fewer measurements compared to conventional methods, while maintaining high fidelity. Evidence shows that adaptive measurement strategies driven by information gain

significantly reduce redundancy and accelerate convergence, particularly in large Hilbert spaces and low-rank states. The study further reveals that reconstruction efficiency does not come at the expense of robustness, as the adaptive framework maintains superior performance under moderate noise conditions. This finding differentiates the research from prior approaches by establishing that scalability and precision in quantum state reconstruction can be achieved simultaneously through intelligent measurement selection rather than exhaustive sampling.

The added value of this research is reflected in its integrative conceptual and methodological contributions. Conceptually, the study advances a unified framework that combines adaptive measurement selection with robust reconstruction algorithms, bridging the gap between information-theoretic optimization and practical quantum state estimation. Methodologically, the use of large-scale simulations across varying dimensions, state structures, and noise conditions provides a comprehensive and systematic evaluation of performance. The integration of Bayesian updating, information gain criteria, and convex optimization distinguishes this approach from existing methods that treat measurement and reconstruction as separate processes, offering a more efficient and application-oriented solution for quantum tomography.

Several limitations should be acknowledged, which also point toward future research directions. The study is primarily based on theoretical modeling and numerical simulations, limiting direct validation in experimental quantum systems. Assumptions regarding noise models and prior distributions may not fully capture the complexity of real-world quantum devices. Future research should focus on experimental implementation of adaptive tomography, integration with machine learning techniques for enhanced measurement selection, and extension to ultra-high-dimensional and multi-partite quantum systems. Further investigation into real-time adaptive protocols and hardware-level optimization is necessary to enable practical deployment in next-generation quantum technologies.

DECLARATION OF AI AND AI ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

During the preparation of this work the author(s) used [NAME TOOL / SERVICE] in order to [REASON]. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

ACKNOWLEDGMENTS

I would like to thank you funding for this research. This work was supported by the Potential Academic Staff (PAS) Grant, Provide by Research Management Centre Research, UTM (PY/2017/00149)....

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.
Author 2: Conceptualization; Data curation; In-vestigation.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Acharya, J. (2025). Pauli Measurements Are Not Optimal for Single-Copy Tomography. *Proceedings of the Annual ACM Symposium on Theory of Computing*, (Query date: 2026-03-30 20:00:42), 718–729. <https://doi.org/10.1145/3717823.3718248>
- Andrade, M. G. D. (2023). On the Characterization of Quantum Flip Stars with Quantum Network Tomography. *Proceedings 2023 IEEE International Conference on Quantum Computing and Engineering Qce 2023, 1*(Query date: 2026-03-30 20:00:42), 1260–1270. <https://doi.org/10.1109/QCE57702.2023.00142>
- Anshu, A. (2022). Distributed Quantum inner product estimation. *Proceedings of the Annual ACM Symposium on Theory of Computing*, (Query date: 2026-03-30 20:00:42), 44–51. <https://doi.org/10.1145/3519935.3519974>
- Bădescu, C. (2024). Improved quantum data analysis. *Theoretics*, 3(Query date: 2026-03-30 20:00:42). <https://doi.org/10.46298/theoretics.24.7>
- Berritta, F. (2025). Efficient Qubit Calibration by Binary-Search Hamiltonian Tracking. *Prx Quantum*, 6(3). <https://doi.org/10.1103/77qg-p68k>
- Bogdanov, Y. I. (2022). High-fidelity tracking of the evolution of multilevel quantum states. *Proceedings of SPIE the International Society for Optical Engineering, 12157*(Query date: 2026-03-30 20:00:42). <https://doi.org/10.1117/12.2624232>
- Chen, S. (2023). When Does Adaptivity Help for Quantum State Learning? *Proceedings Annual IEEE Symposium on Foundations of Computer Science Focs*, (Query date: 2026-03-30 20:00:42), 391–404. <https://doi.org/10.1109/FOCS57990.2023.00029>
- Chen, X. (2024). Adaptive Online Learning of Quantum States. *Quantum*, 8(Query date: 2026-03-30 20:00:42). <https://doi.org/10.22331/q-2024-09-12-1471>
- Cifarelli, V. (2022). Cardiac immune cell infiltration associates with abnormal lipid metabolism. *Frontiers in Cardiovascular Medicine*, 9(Query date: 2026-03-30 20:00:42). <https://doi.org/10.3389/fcvm.2022.948332>
- Cong, S. (2024). Pure State Feedback Switching Control Based on the Online Estimated State for Stochastic Open Quantum Systems. *IEEE Caa Journal of Automatica Sinica*, 11(10), 2166–2178. <https://doi.org/10.1109/JAS.2023.124071>
- Evans, T. J. (2022). Fast Bayesian Tomography of a Two-Qubit Gate Set in Silicon. *Physical Review Applied*, 17(2). <https://doi.org/10.1103/PhysRevApplied.17.024068>
- Farooq, A. (2022). Self-guided quantum state learning for mixed states. *Quantum Information Processing*, 21(7). <https://doi.org/10.1007/s11128-022-03585-8>
- Fernandez, E. J. (2024). Handbook of Adaptive Optics: From Foundations to Applications. In *Handbook of Adaptive Optics from Foundations to Applications* (p. 275). <https://doi.org/10.1201/9781003163671>
- Gier, D. (2025). Intrinsic and Measured Information in Separable Quantum Processes. *Entropy*, 27(6). <https://doi.org/10.3390/e27060599>
- Hwang, H. (2022). Adaptive Quantum Tomography in Weak Measurement with Superconducting Circuits. *Proceedings 2022 IEEE International Conference on Quantum Computing and Engineering Qce 2022*, (Query date: 2026-03-30 20:00:42), 733–735. <https://doi.org/10.1109/QCE53715.2022.00101>

- Hwang, H. (2023). Adaptive quantum tomography in an indistinct measurement system with superconducting circuits. *Physical Review Applied*, 20(6).
<https://doi.org/10.1103/PhysRevApplied.20.064007>
- Lange, H. (2023). Adaptive Quantum State Tomography with Active Learning. *Quantum*, 7(Query date: 2026-03-30 20:00:42). <https://doi.org/10.22331/q-2023-10-09-1129>
- Laskar, M. A. R. (2024). Adaptive Scalpel Scanning Probe Microscopy for Enhanced Volumetric Sensing in Tomographic Analysis. *Advanced Materials Interfaces*, 11(21).
<https://doi.org/10.1002/admi.202400187>
- Li, B. (2023). Optimal single-qubit tomography: Realization of locally optimal measurements on a quantum computer. *Physical Review A*, 108(3).
<https://doi.org/10.1103/PhysRevA.108.032605>
- Lohani, S. (2023). Dimension-adaptive machine learning-based quantum state reconstruction. *Quantum Machine Intelligence*, 5(1). <https://doi.org/10.1007/s42484-022-00088-8>
- Mai, T. T. (2023). An efficient adaptive MCMC algorithm for Pseudo-Bayesian quantum tomography. *Computational Statistics*, 38(2), 827–843. <https://doi.org/10.1007/s00180-022-01264-x>
- Malik, J. (2022). 3D Quantum Cuts for automatic segmentation of porous media in tomography images. *Computers and Geosciences*, 159(Query date: 2026-03-30 20:00:42).
<https://doi.org/10.1016/j.cageo.2021.105017>
- Mansouri, A. (2023). Experimental Demonstration of Neural Adaptive Quantum Tomography for Two-Qubit States. *Laser Science Ls 2023 in Proceedings Frontiers in Optics Laser Science 2023 Fio Ls Part of Frontiers in Optics Laser Science 2023*, (Query date: 2026-03-30 20:00:42).
<https://www.scopus.com/inward/record.uri?partnerID=HzOxMe3b&scp=85215510846&origin=inward>
- Meng, X. (2022). Pure state tomography with adaptive Pauli measurements. *Journal of University of Science and Technology of China*, 52(8). <https://doi.org/10.52396/JUSTC-2022-0037>
- Mondal, S. (2023). A Bayesian quantum state tomography along with adaptive frameworks based on linear minimum mean square error criterion. *New Journal of Physics*, 25(12).
<https://doi.org/10.1088/1367-2630/ad0e49>
- Nakamura, Y. (2024). Adaptive measurement strategy for quantum subspace methods. *New Journal of Physics*, 26(3). <https://doi.org/10.1088/1367-2630/ad2c3b>
- Nayak, A. (2025). Lower Bounds for Learning Quantum States with Single-Copy Measurements. *ACM Transactions on Computation Theory*, 17(1).
<https://doi.org/10.1145/3717450>
- Palanisamy, K. P. (2024). PART IV: Quantum evaluation models: Quantum process tomography and regression. *Quantum Machine Learning A Modern Approach*, (Query date: 2026-03-30 20:00:42), 262–279. <https://doi.org/10.1201/9781003429654-16>
- Šafránek, D. (2023). Quantifying information extraction using generalized quantum measurements. *Physical Review A*, 108(3).
<https://doi.org/10.1103/PhysRevA.108.032413>
- Silva, L. de. (2024). Unraveling devitalization: Its impact on immune response and ectopic bone remodeling from autologous and allogeneic callus mimics. *Stem Cells Translational Medicine*, 13(11), 1086–1100. <https://doi.org/10.1093/stcltm/szae063>

- Sivak, V. V. (2022). Model-Free Quantum Control with Reinforcement Learning. *Physical Review X*, 12(1). <https://doi.org/10.1103/PhysRevX.12.011059>
- Vorbau, R. (2024). Task-based image quality assessment of an intraoperative CBCT for spine surgery compared with conventional CT. *Physica Medica*, 124(Query date: 2026-03-30 20:00:42). <https://doi.org/10.1016/j.ejmp.2024.103426>
- Wang, Y. (2022). Pure State Tomography with Fourier Transformation. *Advanced Quantum Technologies*, 5(8). <https://doi.org/10.1002/qute.202100091>
- Xiao, S. (2022). Optimal and two-step adaptive quantum detector tomography. *Automatica*, 141(Query date: 2026-03-30 20:00:42). <https://doi.org/10.1016/j.automatica.2022.110296>
- Xiao, S. (2023). Quantum state and detector tomography with known rank. *IFAC Papersonline*, 56(2), 5881–5887. <https://doi.org/10.1016/j.ifacol.2023.10.092>
- Xing, Q. (2025). Enhancing photon-counting computed tomography reconstruction via subspace dictionary learning and spatial sparsity regularization. *Quantitative Imaging in Medicine and Surgery*, 15(1), 581–607. <https://doi.org/10.21037/qims-24-1248>
- Yu, Q. (2025). Advances and Applications of Shack–Hartmann Wavefront Sensors (Invited). *Guangxue Xuebao Acta Optica Sinica*, 45(21). <https://doi.org/10.3788/AOS251187>
- Zhan, H. (2025). Experimental benchmarking of quantum state overlap estimation strategies with photonic systems. *Light Science and Applications*, 14(1). <https://doi.org/10.1038/s41377-025-01755-8>
- Zhang, L. (2025). Multi-scale attention encoder-decoder network for low-dose dental CBCT denoising. *Journal of Image and Graphics*, 30(11), 3694–3706. <https://doi.org/10.11834/jig.250055>
- Zhao, Z. (2025). QuFM: Towards Efficient Quantum Link Fidelity Measurements in Quantum Networks. *Proceedings 2025 International Conference on Quantum Communications Networking and Computing Qcnc 2025*, (Query date: 2026-03-30 20:00:42), 516–520. <https://doi.org/10.1109/QCNC64685.2025.00086>
- Zheng, Y. (2024). Characterizing Biphoton Spatial Wave Function Dynamics with Quantum Wavefront Sensing. *Physical Review Letters*, 133(3). <https://doi.org/10.1103/PhysRevLett.133.033602>
- Zhou, S. (2024). Research Progress in Orbital Angular Momentum Recognition for Laser Beams Based on Artificial Intelligence (Invited). *Guangxue Xuebao Acta Optica Sinica*, 44(14). <https://doi.org/10.3788/AOS231987>

Copyright Holder :

© Aram Hakobyan et al. (2026).

First Publication Right :

© Journal of Tecnologia Quantica

This article is under: