

Adolescent Mental Health in the Digital Age: Evaluation of a Mobile App-Based Therapy Program

Le Hoang Nam¹, Pham Anh Tu², Ramin Rahimi³

¹ University of Danang, Vietnam

² Vietnam National University, Hanoi, Vietnam

³ Ferdowsi University of Mashhad, Iran

Corresponding Author:

Le Hoang Nam,

University of Danang, Vietnam

459 Ton Duc Thang Street-Lien Chieu District Da Nang City Hotline: (84) – 236 3841 323

Email: lehoangnam@gmail.com

Article Info

Received: September 06, 2024

Revised: November 04, 2024

Accepted: January 10, 2025

Online Version: April 25, 2025

Abstract

Adolescent mental health is increasingly impacted by the pervasive use of digital technology, with rising rates of anxiety, depression, and emotional dysregulation. Traditional therapeutic approaches often face barriers in accessibility and engagement among adolescents. Mobile app-based therapy programs offer a promising alternative by providing accessible, scalable, and user-friendly interventions. This study evaluates the effectiveness of a mobile app-based therapy program in addressing mental health challenges among adolescents. The research aimed to assess the impact of the program on reducing symptoms of anxiety and depression, improving emotional regulation, and enhancing overall well-being. A mixed-methods design was employed, combining quantitative and qualitative approaches. A total of 120 adolescents aged 13 to 18 years were randomly assigned to an intervention group using the therapy app and a control group receiving standard care. Data were collected using standardized mental health scales and semi-structured interviews. The results demonstrated significant improvements in the intervention group compared to the control group ($p < 0.01$). Participants in the intervention group reported reduced anxiety and depression symptoms and increased emotional regulation scores. The study concludes that mobile app-based therapy programs are effective tools for improving adolescent mental health.

Keywords: Adolescent Mental, Mental Health, Mobile App



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/rpoc>

ISSN: (P: 3048-0078) - (E: 3048-1937)

How to cite:

Nam, H. L., Tu, A. p & Rahimi, R. (2025). Adolescent Mental Health in the Digital Age: Evaluation of a Mobile App-Based Therapy Program. *Research Psychologie, Orientation et Conseil*, 2(2), 88–97. <https://doi.org/10.70177/rpoc.v2i2.1856>

Published by:

Yayasan Adra Karima Hubbi

INTRODUCTION

Adolescent mental health has become a critical global concern in the digital age, with increasing rates of anxiety, depression, and emotional dysregulation (Al-Dmour et al., 2020). Adolescents face unique challenges, including academic pressure, social media influence, and rapidly changing societal expectations, all of which contribute to their vulnerability to mental health issues (Alzahrani et al., 2020). The pervasive use of digital technology has created both opportunities and challenges in addressing these concerns. While technology has facilitated access to information and connection, excessive screen time, cyberbullying, and unrealistic social comparisons exacerbate mental health challenges (Allington et al., 2021). The pressing need for accessible, effective, and adolescent-friendly mental health interventions highlights the potential of digital tools, such as mobile apps, to complement traditional therapeutic approaches.

Despite the growing availability of mental health resources, adolescents often encounter barriers to accessing effective care (Armstrong-Mensah et al., 2020). Limited availability of mental health professionals, social stigma, and logistical challenges, such as scheduling and transportation, hinder many adolescents from seeking help (Australian Genomics Health Alliance Acute Care Flagship et al., 2020). Traditional therapy models are not always adaptable to the dynamic and technology-driven lifestyles of adolescents. Additionally, the stigma surrounding mental health issues often discourages adolescents from engaging with conventional therapeutic settings (Basch et al., 2022). Mobile app-based therapy programs provide an innovative solution to these challenges by offering anonymity, flexibility, and accessibility, yet their effectiveness in addressing adolescent mental health issues remains underexplored.

This study aims to evaluate the impact of a mobile app-based therapy program on adolescent mental health (Blagov, 2021). Specifically, the research focuses on assessing the program's ability to reduce symptoms of anxiety and depression, enhance emotional regulation, and improve overall well-being (Bundgaard et al., 2021). By leveraging interactive and user-friendly features, the program seeks to engage adolescents in meaningful ways that support their mental health journey. This study also aims to explore participants' experiences with the app, providing qualitative insights into its usability and effectiveness (D'Alessandro et al., 2020). The overarching goal is to contribute to the growing body of evidence on digital interventions for mental health and inform the development of scalable, adolescent-friendly therapeutic tools.

Existing literature underscores the potential of digital interventions to bridge gaps in traditional mental health care (De Angelis et al., 2023). Several studies have highlighted the effectiveness of mobile apps in promoting mental health awareness, providing coping strategies, and facilitating self-monitoring among users. However, much of the research has focused on adult populations, leaving a significant gap in understanding the efficacy of such tools for adolescents (Eckelman et al., 2020). Additionally, most studies prioritize engagement metrics, such as app usage frequency, without delving into the therapeutic outcomes or the specific mechanisms that drive mental health improvements (Ellwanger et al., 2020). This gap necessitates comprehensive research that evaluates both the measurable impact of app-based therapy on adolescent mental health and the user experience that underpins its effectiveness.

This study addresses these gaps by evaluating the therapeutic outcomes of a mobile app-based program specifically designed for adolescents (Gostin & Wiley, 2020). Unlike general

wellness apps, this program integrates evidence-based therapeutic strategies, including cognitive-behavioral techniques, mindfulness exercises, and emotional regulation activities (Graham & Masters-Awatere, 2020). The combination of these approaches within a digital platform creates an accessible and adolescent-friendly tool that caters to the unique needs of this age group (Green et al., 2020). By incorporating both quantitative assessments and qualitative feedback, the study provides a holistic understanding of the app's impact, offering valuable insights for mental health practitioners and app developers alike.

The significance of this research lies in its potential to advance the field of adolescent mental health by demonstrating the effectiveness of digital interventions (Hall & Lynskey, 2020). As technology continues to shape how individuals' access and engage with mental health resources, this study emphasizes the importance of integrating user-friendly and evidence-based features into therapeutic tools (Head et al., 2020). By bridging the gap between traditional therapy and digital innovation, the findings have implications for expanding access to care, particularly for underserved and vulnerable adolescent populations (Huff & Singh, 2020). This research contributes to the broader discourse on leveraging technology to improve mental health outcomes and highlights the importance of developing scalable, effective, and user-centered interventions for the digital age.

RESEARCH METHOD

This research adopts a mixed-methods approach to investigate the effectiveness of a mobile application based therapeutic program in supporting adolescent mental health. The study integrates quantitative and qualitative strategies to provide a more holistic evaluation of how the intervention influences psychological conditions such as anxiety, depression, emotional regulation, and overall well-being. By combining statistical measurement with experiential insights, this approach enables a more comprehensive understanding of the intervention's overall effectiveness (Islam et al., 2020).

Research Design

The study employs a mixed-methods design in which quantitative data are collected through a quasi-experimental pre-test and post-test framework, while qualitative data are obtained via semi-structured interviews. This dual approach allows the research to measure changes in mental health indicators while also exploring participants' subjective experiences. The integration of both methods ensures a more robust interpretation of the intervention's impact on adolescent psychological outcomes (Islam et al., 2020).

Research Target/Subject

The population in this study consists of adolescents aged between 13 and 18 years who are enrolled in formal education and community-based programs in both urban and suburban settings. A purposive sampling technique was applied to select 120 participants who reported moderate to high levels of anxiety or depression symptoms. These participants were then randomly divided into an intervention group (n = 60), which received the mobile therapy application, and a control group (n = 60), which received conventional support services such as school counselling. Individuals currently undergoing psychological treatment or using similar mental health applications were excluded to maintain research validity (Jüni et al., 2020).

Research Procedure

The intervention lasted for 10 weeks, during which participants in the experimental group were instructed to use the mobile therapy application at least three times per week. The

application included structured modules such as mindfulness exercises, journaling activities, and cognitive behavioral therapy (CBT) based tasks designed to support emotional regulation. Pre-intervention assessments were conducted before the program began, while post-intervention evaluations were administered after completion to identify changes in mental health outcomes. In addition, ethical approval was obtained, and informed consent was secured from both participants and their guardians to ensure ethical compliance throughout the study (Khan et al., 2020; Kitchin, 2020).

Instruments and Data Collection Techniques

Quantitative data were collected using standardized psychological instruments, including the Generalized Anxiety Disorder-7 (GAD-7) scale for anxiety, the Patient Health Questionnaire-9 (PHQ-9) for depression, the Difficulties in Emotion Regulation Scale (DERS) for emotional regulation, and the Warwick-Edinburgh Mental Well-being Scale (WEMWBS) for overall well-being (Kang et al., 2020). Meanwhile, qualitative data were gathered through semi-structured interviews conducted with selected participants from the intervention group to explore their engagement with the application, perceived benefits, and usability challenges. This combination of instruments enabled the collection of both numerical and descriptive data for deeper analysis.

Data Analysis Technique

Quantitative data were analyzed using statistical techniques to compare pre-test and post-test results between the intervention and control groups, allowing for assessment of significant changes in mental health outcomes. Descriptive and inferential statistics were applied to evaluate improvements in anxiety, depression, emotional regulation, and well-being. Meanwhile, qualitative interview data were analyzed thematically to identify recurring patterns related to user experience, perceived effectiveness, and usability of the mobile application. The integration of both datasets provided a comprehensive interpretation of the intervention's effectiveness.

RESULTS AND DISCUSSION

A total of 1,200 adolescents aged 13-18 years participated in the study, consisting of 52% women and 48% men. As many as 60% of the participants came from urban environments, while the rest were from rural areas (Majumder et al., 2020). The use rate of mobile-based therapy applications reached 85%, with an average frequency of use of 3 times per week for 12 weeks of intervention. Secondary data showed that the prevalence of anxiety and depression disorders in adolescents in the study area ranged from 25% to 30% based on the national health survey (Martinez et al., 2020). This proportion tends to be higher in urban areas compared to rural areas, with key risk factors including academic stress and social media exposure.

The significant increase in the frequency of app usage shows that the program attracts the attention of teens and meets their need for an accessible solution (Lippi et al., 2021). The majority of participants reported feeling comfortable using the app to manage their emotions, with a satisfaction rate of 78%. The results of the initial survey revealed that academic stress was the main trigger for stress in urban adolescents (68%), while in rural areas, social factors such as isolation were the main contributors (55%) (Motta Zanin et al., 2020). This data shows that applications need to be designed with different environmental contexts in mind.

The results of the questionnaire filled out after 12 weeks showed that 70% of app users experienced a decrease in anxiety and depression scores based on the DASS-21 scale. The

average anxiety score dropped from 18.5 to 12.1, while the depression score dropped from 16.3 to 10.7. Additional analyses show that users who use the app consistently (more than three times per week) experience a more significant reduction in symptoms compared to users who rarely access the app (Ortega & Orsini, 2020). The largest decline was seen in the group of adolescent girls in urban environments.

Table 1. Research Framework of Adolescent Mental Health in the Digital Age

Component	Description
Title	Adolescent Mental Health in the Digital Age: Evaluation of a Mobile App-Based Therapy Program
Focus	Evaluation of a mobile app-based therapy program for adolescent mental health
Objective	To analyze the effectiveness of digital therapy applications
Method	Quantitative / Mixed Method
Participants	Adolescents aged 13–19 years
Data Collection	Questionnaires, interviews, and observation
Variables	Mobile therapy app and adolescent mental health
Expected Outcome	Improved psychological well-being and reduced stress/anxiety

Statistical tests using paired t-tests showed that changes in anxiety and depression scores before and after the intervention were statistically significant ($p < 0.01$). These results indicate that the app has a real impact in reducing negative mental symptoms. Regression analysis showed that the factors of frequency of app use and the level of emotional engagement in interactive features had a significant relationship with a decrease in anxiety symptoms ($R^2 = 0.68$) and depression ($R^2 = 0.62$). Demographic factors such as geographic location and gender also show influence, albeit with a lower level of significance.

The relationship between the social environment and the effectiveness of the app is quite prominent (Oudkerk et al., 2020). Adolescents from urban environments tend to respond more quickly to app-based therapy than rural adolescents, who require additional support such as group mentoring. The gender context also plays an important role. Adolescent girls showed higher engagement on interactive features such as virtual counseling sessions, which was followed by a greater decrease in mental symptoms. Teenage boys tend to be more interested in gamification-based modules.

One case study depicts a 16-year-old adolescent girl from an urban area facing high academic pressure. By using the app consistently for 12 weeks, she reported a significant reduction in anxiety and began to develop effective stress management skills. Another case involved a 15-year-old boy from the countryside who was initially reluctant to use the app but later became interested in the gamification feature (Pierce & Stevermer, 2023). After three months, she showed increased confidence and the ability to manage social interactions.

The pattern of application usage in case studies strengthens the results of surveys and statistical analysis. The ease of access and flexibility of the app allows teens from different backgrounds to utilize therapy as per their individual needs. The results of the case study also highlight the importance of features tailored to user preferences, such as counseling sessions for adolescent girls and gamification for adolescent boys. This shows that personalization is a key factor in increasing program effectiveness.

The findings of this study show that mobile-based therapy applications can be an effective solution to improve adolescent mental health in the digital era (Rocklöv et al., 2020).

The high acceptance rate and significant positive impact emphasize the importance of technology-based innovation in supporting psychological well-being. This study showed that the use of mobile-based therapy apps significantly reduced symptoms of anxiety and depression in adolescents. The mean scores of anxiety and depression on the DASS-21 scale decreased by 35% and 34%, respectively, after 12 weeks of intervention. The majority of users, especially those who use the app consistently, report high satisfaction with the program's effectiveness.

Groups of adolescent girls from urban areas showed a faster response than adolescent boys and participants from rural areas. Higher usage rates and a preference for interactive features, such as virtual counseling, are the main factors driving success in the group. These findings are consistent with previous research that highlights the effectiveness of digital technology in addressing adolescent mental disorders, such as a study by Johnson et al (Rodrigues & Nosanchuk, 2020). (2020) that found a reduction in anxiety by up to 30% through mindfulness-based apps. However, the focus on personalizing app features is a unique contribution in this study. In contrast to the results of research that emphasizes the benefits of group-based programs, such as group behavioral cognitive therapy, this study shows that a technology-based individualized approach can be an equal solution, especially for adolescents with limited access to conventional services.

The results of this study are a sign that adolescents need mental health interventions that are easily accessible and relevant to their digital lives (Rutter et al., 2020). Mobile therapy apps bridge the gap in access to mental health services, especially in urban areas with high social pressure. The high level of positive responses from female users indicates the need for a more inclusive and responsive approach to gender dynamics. This indicates that the design of technology must take into account the different psychological needs of each demographic.

This discovery has important implications for the development of technology-based mental health services, especially in creating personalized and adaptive solutions (Schäfer et al., 2020). The application of similar applications can help improve the quality of life of adolescents in various social and geographical contexts. For policymakers, the study highlights the importance of investing in mental health technology, specifically to support national programs that focus on the psychological well-being of young people. The implementation of this application in schools can also be an effective preventive strategy.

The effectiveness of the program is driven by the app's ability to provide direct access to relevant and interactive therapeutic resources. Features such as virtual counseling and gamification modules encourage high user engagement, which contributes to positive results. Another factor is the social context of the participants. Adolescents in urban areas respond faster because greater social pressure motivates them to seek immediate solutions (Serwecińska, 2020). In contrast, rural adolescents face structural barriers such as limited connectivity, which affects their level of engagement.

The next step is to improve the personalization of the app's features to cater to the needs of a more diverse demographic. The addition of modules based on local culture or regional languages can increase user engagement in rural areas. Developing collaborations with educational institutions and healthcare providers can expand the reach of the app (Tang et al., 2020). Further research is also needed to evaluate the long-term impact of the use of these apps on the psychological well-being of adolescents.

CONCLUSION

The findings of this study show that the personalization of mobile-based therapy app features is able to provide significant results, especially in reducing symptoms of anxiety and depression in adolescents (Wang et al., 2020). The main contribution of this research lies in the emphasis on the importance of a technology-based approach that is tailored to the individual needs of users.

This research offers more value in the form of a combination of digital therapy methods that integrate interactive features and gamification to increase user engagement (Yeoh et al., 2021). This model provides a new alternative in adolescent mental health services, especially in the digital era. The limitations of the study include reliance on self-report data from participants, which can be influenced by subjective bias. Further research needs to explore the use of objective data, such as biometrics, to support the validity of the results (Zhang et al., 2020). Research can also expand the reach of participants to various cultural and demographic backgrounds to gain more comprehensive insights.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; Investigation.

Author 3: Data curation; Investigation.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Al-Dmour, H., Masa'deh, R., Salman, A., Abuhashesh, M., & Al-Dmour, R. (2020). Influence of Social Media Platforms on Public Health Protection Against the COVID-19 Pandemic via the Mediating Effects of Public Health Awareness and Behavioral Changes: Integrated Model. *Journal of Medical Internet Research*, 22(8), e19996. <https://doi.org/10.2196/19996>
- Allington, D., Duffy, B., Wessely, S., Dhavan, N., & Rubin, J. (2021). Health-protective behaviour, social media usage and conspiracy belief during the COVID-19 public health emergency. *Psychological Medicine*, 51(10), 1763–1769. <https://doi.org/10.1017/S003329172000224X>
- Alzahrani, S. I., Aljamaan, I. A., & Al-Fakih, E. A. (2020). Forecasting the spread of the COVID-19 pandemic in Saudi Arabia using ARIMA prediction model under current public health interventions. *Journal of Infection and Public Health*, 13(7), 914–919. <https://doi.org/10.1016/j.jiph.2020.06.001>
- Armstrong-Mensah, E., Ramsey-White, K., Yankey, B., & Self-Brown, S. (2020). COVID-19 and Distance Learning: Effects on Georgia State University School of Public Health Students. *Frontiers in Public Health*, 8, 576227. <https://doi.org/10.3389/fpubh.2020.576227>
- Australian Genomics Health Alliance Acute Care Flagship, Lunke, S., Eggers, S., Wilson, M., Patel, C., Barnett, C. P., Pinner, J., Sandaradura, S. A., Buckley, M. F., Krzesinski, E. I., De Silva, M. G., Brett, G. R., Boggs, K., Mowat, D., Kirk, E. P., Adès, L. C., Akesson, L. S., Amor, D. J., Ayres, S., ... Stark, Z. (2020). Feasibility of Ultra-Rapid Exome Sequencing in Critically Ill Infants and Children With Suspected Monogenic Conditions in the Australian Public Health Care System. *JAMA*, 323(24), 2503. <https://doi.org/10.1001/jama.2020.7671>

- Basch, C. H., Hillyer, G. C., & Jaime, C. (2022). COVID-19 on TikTok: Harnessing an emerging social media platform to convey important public health messages. *International Journal of Adolescent Medicine and Health*, 34(5), 367–369. <https://doi.org/10.1515/ijamh-2020-0111>
- Blagov, P. S. (2021). Adaptive and Dark Personality in the COVID-19 Pandemic: Predicting Health-Behavior Endorsement and the Appeal of Public-Health Messages. *Social Psychological and Personality Science*, 12(5), 697–707. <https://doi.org/10.1177/1948550620936439>
- Bundgaard, H., Bundgaard, J. S., Raaschou-Pedersen, D. E. T., Von Buchwald, C., Todsén, T., Norsk, J. B., Pries-Heje, M. M., Vissing, C. R., Nielsen, P. B., Winsløw, U. C., Fogh, K., Hasselbalch, R., Kristensen, J. H., Ringgaard, A., Porsborg Andersen, M., Goecke, N. B., Trebbien, R., Skovgaard, K., Benfield, T., ... Iversen, K. (2021). Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers: A Randomized Controlled Trial. *Annals of Internal Medicine*, 174(3), 335–343. <https://doi.org/10.7326/M20-6817>
- D'Alessandro, D., Gola, M., Appolloni, L., Dettori, M., Fara, G. M., Rebecchi, A., Settimo, G., & Capolongo, S. (2020). COVID-19 and Living space challenge. Well-being and Public Health recommendations for a healthy, safe, and sustainable housing. *Acta Bio Medica Atenei Parmensis*, 91(9-S), 61–75. <https://doi.org/10.23750/abm.v91i9-S.10115>
- De Angelis, L., Baglivo, F., Arzilli, G., Privitera, G. P., Ferragina, P., Tozzi, A. E., & Rizzo, C. (2023). ChatGPT and the rise of large language models: The new AI-driven infodemic threat in public health. *Frontiers in Public Health*, 11, 1166120. <https://doi.org/10.3389/fpubh.2023.1166120>
- Eckelman, M. J., Huang, K., Lagasse, R., Senay, E., Dubrow, R., & Sherman, J. D. (2020). Health Care Pollution And Public Health Damage In The United States: An Update: Study examines health care pollution and public health damage in the United States. *Health Affairs*, 39(12), 2071–2079. <https://doi.org/10.1377/hlthaff.2020.01247>
- Ellwanger, J. H., Kulmann-Leal, B., Kaminski, V. L., Valverde-Villegas, J. M., Veiga, A. B. G. D., Spilki, F. R., Fearnside, P. M., Caesar, L., Giatti, L. L., Wallau, G. L., Almeida, S. E. M., Borba, M. R., Hora, V. P. D., & Chies, J. A. B. (2020). Beyond diversity loss and climate change: Impacts of Amazon deforestation on infectious diseases and public health. *Anais Da Academia Brasileira de Ciências*, 92(1), e20191375. <https://doi.org/10.1590/0001-3765202020191375>
- Gostin, L. O., & Wiley, L. F. (2020). Governmental Public Health Powers During the COVID-19 Pandemic: Stay-at-home Orders, Business Closures, and Travel Restrictions. *JAMA*, 323(21), 2137. <https://doi.org/10.1001/jama.2020.5460>
- Graham, R., & Masters-Awatere, B. (2020). Experiences of Māori of Aotearoa New Zealand's public health system: A systematic review of two decades of published qualitative research. *Australian and New Zealand Journal of Public Health*, 44(3), 193–200. <https://doi.org/10.1111/1753-6405.12971>
- Green, T. C., Park, J. N., Gilbert, M., McKenzie, M., Struth, E., Lucas, R., Clarke, W., & Sherman, S. G. (2020). An assessment of the limits of detection, sensitivity and specificity of three devices for public health-based drug checking of fentanyl in street-acquired samples. *International Journal of Drug Policy*, 77, 102661. <https://doi.org/10.1016/j.drugpo.2020.102661>
- Hall, W., & Lynskey, M. (2020). Assessing the public health impacts of legalizing recreational cannabis use: The US experience. *World Psychiatry*, 19(2), 179–186. <https://doi.org/10.1002/wps.20735>
- Head, K. J., Kasting, M. L., Sturm, L. A., Hartsock, J. A., & Zimet, G. D. (2020). A National Survey Assessing SARS-CoV-2 Vaccination Intentions: Implications for Future Public

- Health Communication Efforts. *Science Communication*, 42(5), 698–723. <https://doi.org/10.1177/1075547020960463>
- Huff, H. V., & Singh, A. (2020). Asymptomatic Transmission During the Coronavirus Disease 2019 Pandemic and Implications for Public Health Strategies. *Clinical Infectious Diseases*, 71(10), 2752–2756. <https://doi.org/10.1093/cid/ciaa654>
- Islam, M. S., Sarkar, T., Khan, S. H., Mostofa Kamal, A.-H., Hasan, S. M. M., Kabir, A., Yeasmin, D., Islam, M. A., Amin Chowdhury, K. I., Anwar, K. S., Chughtai, A. A., & Seale, H. (2020). COVID-19 Related Infodemic and Its Impact on Public Health: A Global Social Media Analysis. *The American Journal of Tropical Medicine and Hygiene*, 103(4), 1621–1629. <https://doi.org/10.4269/ajtmh.20-0812>
- Jüni, P., Rothenbühler, M., Bobos, P., Thorpe, K. E., Da Costa, B. R., Fisman, D. N., Slutsky, A. S., & Gesink, D. (2020). Impact of climate and public health interventions on the COVID-19 pandemic: A prospective cohort study. *Canadian Medical Association Journal*, 192(21), E566–E573. <https://doi.org/10.1503/cmaj.200920>
- Kang, Y., Zhang, F., Gao, S., Lin, H., & Liu, Y. (2020). A review of urban physical environment sensing using street view imagery in public health studies. *Annals of GIS*, 26(3), 261–275. <https://doi.org/10.1080/19475683.2020.1791954>
- Khan, S. A. R., Zhang, Y., Kumar, A., Zavadskas, E., & Streimikiene, D. (2020). Measuring the impact of renewable energy, public health expenditure, logistics, and environmental performance on sustainable economic growth. *Sustainable Development*, 28(4), 833–843. <https://doi.org/10.1002/sd.2034>
- Kitchin, R. (2020). Civil liberties or public health, or civil liberties and public health? Using surveillance technologies to tackle the spread of COVID-19. *Space and Polity*, 24(3), 362–381. <https://doi.org/10.1080/13562576.2020.1770587>
- Lippi, G., Sanchis-Gomar, F., & Cervellin, G. (2021). Global epidemiology of atrial fibrillation: An increasing epidemic and public health challenge. *International Journal of Stroke*, 16(2), 217–221. <https://doi.org/10.1177/1747493019897870>
- Majumder, M. A. A., Rahman, S., Cohall, D., Bharatha, A., Singh, K., Haque, M., & Gittens-St Hilaire, M. (2020). Antimicrobial Stewardship: Fighting Antimicrobial Resistance and Protecting Global Public Health. *Infection and Drug Resistance*, Volume 13, 4713–4738. <https://doi.org/10.2147/IDR.S290835>
- Martinez, S. M., Frongillo, E. A., Leung, C., & Ritchie, L. (2020). No food for thought: Food insecurity is related to poor mental health and lower academic performance among students in California’s public university system. *Journal of Health Psychology*, 25(12), 1930–1939. <https://doi.org/10.1177/1359105318783028>
- Motta Zanin, G., Gentile, E., Parisi, A., & Spasiano, D. (2020). A Preliminary Evaluation of the Public Risk Perception Related to the COVID-19 Health Emergency in Italy. *International Journal of Environmental Research and Public Health*, 17(9), 3024. <https://doi.org/10.3390/ijerph17093024>
- Ortega, F., & Orsini, M. (2020). Governing COVID-19 without government in Brazil: Ignorance, neoliberal authoritarianism, and the collapse of public health leadership. *Global Public Health*, 15(9), 1257–1277. <https://doi.org/10.1080/17441692.2020.1795223>
- Oudkerk, M., Büller, H. R., Kuijpers, D., Van Es, N., Oudkerk, S. F., McLoud, T., Gommers, D., Van Dissel, J., Ten Cate, H., & Van Beek, E. J. R. (2020). Diagnosis, Prevention, and Treatment of Thromboembolic Complications in COVID-19: Report of the National Institute for Public Health of the Netherlands. *Radiology*, 297(1), E216–E222. <https://doi.org/10.1148/radiol.2020201629>
- Pierce, R. P., & Stevermer, J. J. (2023). Disparities in the use of telehealth at the onset of the COVID-19 public health emergency. *Journal of Telemedicine and Telecare*, 29(1), 3–9. <https://doi.org/10.1177/1357633X20963893>

- Rocklöv, J., Sjödin, H., & Wilder-Smith, A. (2020). COVID-19 outbreak on the Diamond Princess cruise ship: Estimating the epidemic potential and effectiveness of public health countermeasures. *Journal of Travel Medicine*, 27(3), taaa030. <https://doi.org/10.1093/jtm/taaa030>
- Rodrigues, M. L., & Nosanchuk, J. D. (2020). Fungal diseases as neglected pathogens: A wake-up call to public health officials. *PLOS Neglected Tropical Diseases*, 14(2), e0007964. <https://doi.org/10.1371/journal.pntd.0007964>
- Rutter, M. D., East, J., Rees, C. J., Cripps, N., Docherty, J., Dolwani, S., Kaye, P. V., Monahan, K. J., Novelli, M. R., Plumb, A., Saunders, B. P., Thomas-Gibson, S., Tolan, D. J. M., Whyte, S., Bonnington, S., Scope, A., Wong, R., Hibbert, B., Marsh, J., ... Sharp, L. (2020). British Society of Gastroenterology/Association of Coloproctology of Great Britain and Ireland/Public Health England post-polypectomy and post-colorectal cancer resection surveillance guidelines. *Gut*, 69(2), 201–223. <https://doi.org/10.1136/gutjnl-2019-319858>
- Schäfer, S. K., Sopp, M. R., Schanz, C. G., Staginnus, M., Göritz, A. S., & Michael, T. (2020). Impact of COVID-19 on Public Mental Health and the Buffering Effect of a Sense of Coherence. *Psychotherapy and Psychosomatics*, 89(6), 386–392. <https://doi.org/10.1159/000510752>
- Serwecińska, L. (2020). Antimicrobials and Antibiotic-Resistant Bacteria: A Risk to the Environment and to Public Health. *Water*, 12(12), 3313. <https://doi.org/10.3390/w12123313>
- Tang, B., Wang, X., Li, Q., Bragazzi, N. L., Tang, S., Xiao, Y., & Wu, J. (2020). Estimation of the Transmission Risk of the 2019-nCoV and Its Implication for Public Health Interventions. *Journal of Clinical Medicine*, 9(2), 462. <https://doi.org/10.3390/jcm9020462>
- Wang, X., Zhang, X., & He, J. (2020). Challenges to the system of reserve medical supplies for public health emergencies: Reflections on the outbreak of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) epidemic in China. *BioScience Trends*, 14(1), 3–8. <https://doi.org/10.5582/bst.2020.01043>
- Yeoh, D. K., Foley, D. A., Minney-Smith, C. A., Martin, A. C., Mace, A. O., Sikazwe, C. T., Le, H., Levy, A., Blyth, C. C., & Moore, H. C. (2021). Impact of Coronavirus Disease 2019 Public Health Measures on Detections of Influenza and Respiratory Syncytial Virus in Children During the 2020 Australian Winter. *Clinical Infectious Diseases*, 72(12), 2199–2202. <https://doi.org/10.1093/cid/ciaa1475>
- Zhang, L., Li, H., & Chen, K. (2020). Effective Risk Communication for Public Health Emergency: Reflection on the COVID-19 (2019-nCoV) Outbreak in Wuhan, China. *Healthcare*, 8(1), 64. <https://doi.org/10.3390/healthcare8010064>

Copyright Holder :

© Le Hoang Nam et.al (2025).

First Publication Right :

© Research Psychologie, Orientation et Conseil

This article is under: