OPEN ACCESS
Research of Scientia Naturalis
Vol. 3 No. 5. October 2026, pp. 447-464
DOI. 10.70177/scientia.v3i5.4480
Research Article

DYNAMIC MODELING OF ECOSYSTEM SERVICES UNDER RAPID CLIMATE CHANGE SCENARIOS TO INFORM POLICY DECISIONS FOR BIODIVERSITY CONSERVATION

Achmad Agus Salim[1], Sota Yamamoto[2], and Muhammad Yassir[3]
[1] nstitut Teknologi Sepuluh Nopember, Indonesia
[2] Osaka University, Japan
[3] Universitas Gunung Leuser, Indonesia

Corresponding Author:

Achmad Agus Salim,
Department of Environmental Engineering, Faculty of Civil Engineering, Planning, and Geosciences, Institut Teknologi Sepuluh Nopember.
Jl. Raya ITS, Sukolilo Surabaya 60111, Indonesia
Email: achmadasalim@gmail.com

Article Info

Received: April 6, 2026
Revised: July 9, 2026
Accepted: September 11, 2026
Online Version: October 27, 2026

Abstract

Rapid climate change has intensified pressures on ecological systems, threatening biodiversity stability and reducing the capacity of ecosystems to provide essential services for human and environmental sustainability. Dynamic modeling offers a promising approach to predict ecosystem responses by integrating climate scenarios, biodiversity patterns, and ecological processes to support adaptive conservation strategies. This study aimed to develop a dynamic modeling framework for evaluating ecosystem service changes under different climate change scenarios and generating scientific evidence for biodiversity conservation policy decisions. A quantitative ecological modeling approach was employed by integrating historical climate data, biodiversity indicators, remote sensing observations, and ecosystem service assessments from forest, freshwater, coastal, and agricultural ecosystems. Data were analyzed through scenario simulation, spatial modeling, regression analysis, and sensitivity evaluation to identify ecological vulnerabilities and adaptation pathways. The findings revealed that climate change significantly influenced ecosystem service capacity, with severe scenarios producing substantial reductions in biodiversity support, habitat stability, and ecological productivity. The results demonstrated that biodiversity richness and habitat connectivity played crucial roles in strengthening ecosystem resilience. The study concludes that dynamic modeling provides an effective decision-support framework for developing proactive conservation policies, enabling policymakers to anticipate ecological risks and implement adaptive strategies for maintaining biodiversity and ecosystem services under future climate uncertainties.

Keywords: Biodiversity Conservation, Climate Change, Dynamic Modeling

© 2026 by the author(s)
This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (https://creativecommons.org/licenses/by-sa/4.0/).

Journal Homepage https://research.adra.ac.id/index.php/scientia
How to cite: Salim, A. A., Yamamoto, S., & Yassir, M. (2026). Dynamic Modeling of Ecosystem Services Under Rapid Climate Change Scenarios to Inform Policy Decisions for Biodiversity Conservation. *Research of Scientia Naturalis*, *3*(5), 447–464. https://doi.org/10.70177/scientia.v3i5.4480
Published by: Yayasan Adra Karima Hubbi

INTRODUCTION

Climate change has emerged as one of the most significant environmental challenges affecting ecological systems, biodiversity stability, and human well-being across the globe (Ghasemi et al., 2025). Increasing temperatures, changing precipitation patterns, extreme weather events, sea-level rise, and ecosystem disturbances have altered the functioning of natural systems and threatened the availability of essential ecosystem services (Elleaume et al., 2025; Grajales Noreña et al., 2024). Ecosystem services, including carbon regulation, water purification, pollination, soil fertility maintenance, and biodiversity support, represent critical connections between ecological processes and human societies (Hulme et al., 2025; Pyron et al., 2024). The accelerating pace of climate change requires improved understanding of how ecosystems respond dynamically under different environmental scenarios to support effective conservation strategies and sustainable policy development (Umwali et al., 2025).

Biodiversity conservation has increasingly shifted from static protection approaches toward adaptive management frameworks capable of responding to rapidly changing ecological conditions (Liu, Wang, et al., 2024). Traditional conservation planning often relies on historical ecological data and fixed management strategies that may not adequately address future uncertainties caused by climate change (Rasmus et al., 2024). Ecosystems operate through complex interactions among species, environmental variables, and human activities, making their future trajectories difficult to predict using conventional assessment methods. Dynamic modeling approaches provide opportunities to simulate ecosystem responses, evaluate potential climate scenarios, and identify management strategies that enhance ecological resilience (Aghaloo & Sharifi, 2025; Zhi et al., 2025).

The integration of dynamic modeling with ecosystem service assessment represents an important advancement in environmental science and policy development (W. Yang et al., 2025). Computational models enable researchers to analyze interactions among climate variables, biodiversity patterns, ecosystem functions, and human pressures over time. These approaches provide valuable insights into possible ecological futures and allow policymakers to evaluate conservation interventions before implementation (G.-Z. Lin & Chiang, 2025; F. Wang et al., 2024). Developing reliable dynamic models under rapid climate change scenarios is therefore essential for informing evidence-based decisions aimed at maintaining biodiversity and sustaining ecosystem services (Lathouwers et al., 2025; Weng et al., 2024).

The increasing intensity of climate change creates substantial uncertainty regarding the future capacity of ecosystems to provide essential services and maintain biodiversity (Liu, Huang, et al., 2024). Many ecosystems are experiencing rapid environmental changes that exceed historical patterns, resulting in habitat degradation, species displacement, ecosystem fragmentation, and declining ecological functions (Liu et al., 2025; L. Wang et al., 2024). Existing conservation strategies often struggle to address these challenges because they are based on limited projections or short-term observations. The inability to accurately predict ecosystem responses under different climate scenarios creates difficulties for policymakers seeking to develop effective conservation strategies (González Hernández et al., 2023; Kwak & Chen, 2025).

Current ecosystem service assessments frequently rely on descriptive evaluations or spatial analyses that provide important information about existing ecological conditions but offer limited understanding of future dynamics (Bizama et al., 2025; Pintado & Knoke, 2025). Ecosystem processes are influenced by multiple interacting factors, including climate variability, land-use change, species interactions, and human interventions (Z. Lin et al., 2025). These complex relationships require analytical frameworks capable of capturing temporal changes and feedback mechanisms within ecological systems. The absence of integrated dynamic models reduces the ability to anticipate emerging biodiversity risks and design proactive conservation measures (Rieder et al., 2025; Tan et al., 2025).

The central problem addressed in this research concerns the limited availability of comprehensive modeling approaches that connect climate change scenarios, ecosystem service dynamics, and biodiversity conservation decision-making. Existing models often focus on specific ecological components without fully integrating multiple ecosystem functions and policy considerations. A more adaptive and predictive approach is required to understand how ecosystems may transform under future climate conditions and how conservation policies can respond effectively. This research addresses the need for dynamic modeling frameworks that support long-term ecological planning and informed environmental governance (Bai et al., 2025; F. Lin et al., 2025).

The primary objective of this study is to develop a dynamic modeling framework for assessing changes in ecosystem services under rapid climate change scenarios. The research aims to simulate ecological responses by integrating climate projections, biodiversity indicators, ecosystem functions, and environmental pressures. The study seeks to identify potential changes in ecosystem service availability and determine how different climate conditions may influence ecological stability over time (Y. Wang & Liu, 2025; S. Yang et al., 2024).

The second objective is to evaluate the relationship between climate-driven ecosystem changes and biodiversity conservation outcomes. The research examines how variations in temperature, precipitation, habitat conditions, and human activities influence species distribution, ecosystem functionality, and ecological resilience. The study aims to identify critical ecological factors that determine vulnerability and adaptive capacity within changing environmental systems.

The third objective is to provide scientific insights that support evidence-based policy decisions for biodiversity conservation. The research seeks to translate modeling outputs into practical recommendations for conservation planning, ecosystem management, and climate adaptation strategies. The expected contribution is the development of a predictive framework that enables policymakers to evaluate future risks, prioritize conservation actions, and improve ecological decision-making under uncertain climate conditions.

Existing research on ecosystem services and climate change has produced extensive knowledge regarding ecological impacts, biodiversity loss, and environmental vulnerability. Previous studies have demonstrated that climate change significantly influences ecosystem functions and species distributions across various landscapes (de Araujo Fonseca & da Cunha Bustamante, 2025; Strange et al., 2024). However, many existing assessments remain focused on current conditions or isolated environmental indicators rather than examining long-term ecosystem trajectories under multiple climate scenarios. The limited integration of dynamic simulation approaches restricts understanding of how ecosystem services may evolve in response to future environmental changes (Banerjee et al., 2025; Wu et al., 2024).

Research on ecological modeling has introduced various approaches, including species distribution models, climate impact assessments, and ecosystem simulations. These methods have contributed valuable information regarding specific ecological processes but often lack comprehensive integration between biodiversity dynamics, ecosystem services, and policy applications. Many models provide technical predictions without sufficiently addressing how results can inform practical conservation decisions (G. Lin et al., 2025; Yu & Ma, 2025). This limitation creates a disconnect between scientific modeling outcomes and environmental governance requirements.

Another significant research gap concerns the limited application of dynamic models that incorporate feedback relationships between ecological systems and human decision-making processes. Ecosystem services are influenced not only by climate variables but also by land management practices, conservation interventions, and policy choices. Existing models frequently underestimate the importance of these interactions when predicting ecological futures. This study addresses the gap by developing an integrated modeling approach that

connects climate scenarios, ecosystem service changes, biodiversity conservation priorities, and policy decision-making processes.

The novelty of this research lies in the development of an integrated dynamic modeling framework that combines ecosystem service evaluation, climate change scenarios, biodiversity analysis, and conservation policy considerations. Previous approaches have often examined these components separately, limiting their ability to represent the complexity of real-world ecological systems. This study contributes a holistic perspective by treating ecosystems as dynamic networks where environmental changes, biological responses, and management decisions interact continuously.

The conceptual contribution of this research involves advancing understanding of ecosystem services as evolving systems rather than fixed ecological benefits. The study emphasizes that ecosystem services must be evaluated through temporal and adaptive perspectives because climate change continuously modifies ecological conditions. The integration of dynamic modeling provides a more comprehensive approach for examining resilience, vulnerability, and potential transformation pathways within ecosystems. This perspective contributes to broader discussions on climate adaptation, biodiversity conservation, and sustainable environmental management.

The justification for conducting this research is based on the urgent need for predictive tools that support conservation decisions in an era of rapid environmental change. Policymakers require reliable scientific evidence to allocate conservation resources, prioritize vulnerable ecosystems, and develop adaptive management strategies. Dynamic modeling approaches can reduce uncertainty by providing simulations of possible ecological futures and evaluating alternative policy responses. This research is therefore significant for advancing environmental science while supporting practical efforts to protect biodiversity and maintain ecosystem services under accelerating climate pressures.

RESEARCH METHOD

Research Design

This study employed a quantitative ecological modeling design integrated with spatial analysis and scenario-based simulation to evaluate the dynamics of ecosystem services under rapid climate change conditions. The research adopted an interdisciplinary approach combining ecological science, climate modeling, biodiversity assessment, and environmental policy analysis. The design was developed to examine how variations in climate parameters, land-use changes, and ecological interactions influence the capacity of ecosystems to provide essential services over time. Dynamic modeling was selected because ecosystem responses to climate change involve complex interactions, feedback mechanisms, and temporal variations that cannot be adequately explained through static assessment approaches (Abedin et al., 2025; Y.-P. Lin et al., 2025).

The research framework was based on an ecosystem service dynamics model that integrated climate scenarios, biodiversity indicators, ecological processes, and human-driven environmental pressures. The model incorporated key ecosystem services, including carbon sequestration, water regulation, habitat provision, soil conservation, and biodiversity maintenance. Climate projections were simulated under different future scenarios to evaluate potential ecological changes and identify vulnerability patterns. The modeling framework enabled the assessment of ecosystem responses under alternative environmental conditions and supported the development of evidence-based recommendations for biodiversity conservation policies.

The analytical design combined empirical ecological data with computational simulations to generate predictive insights regarding future ecosystem trajectories. The study applied scenario-based modeling techniques to compare ecosystem responses under different climate

change pathways. Model outputs were evaluated through ecological indicators, including species distribution changes, ecosystem productivity variations, service availability, and resilience capacity. The integration of ecological modeling and policy analysis allowed the research to move beyond impact assessment by identifying potential management strategies that could strengthen ecosystem adaptation under uncertain climate conditions.

Research Target/Subject

The population of this study consisted of ecological systems and biodiversity components within selected representative landscapes experiencing significant climate change pressures. The research focused on ecosystems that provide critical services for ecological stability and human well-being, including forest ecosystems, freshwater systems, coastal environments, and agricultural landscapes. These ecosystems were selected because they represent highly sensitive ecological areas where climate variability, biodiversity changes, and human activities interact strongly (Bazzato et al., 2024; Bianconi et al., 2025).

The study sample was determined using purposive sampling based on ecological importance, vulnerability to climate change, availability of long-term environmental data, and relevance for conservation policy development. Selected ecosystems represented different climatic conditions, biodiversity characteristics, and management contexts to ensure comprehensive evaluation of ecosystem service responses. Sampling criteria included ecosystem diversity, availability of biodiversity records, historical climate information, land-use transformation data, and existing conservation initiatives.

The research analyzed ecological datasets obtained from multiple ecosystem locations representing different environmental conditions. The dataset consisted of climate records from 30 years of historical observations, biodiversity information covering approximately 500 species indicators, land-use change data from satellite observations, and ecosystem service measurements from ecological monitoring programs. The selected samples provided sufficient variation to evaluate how different ecosystems respond to climate scenarios and how conservation interventions may influence future ecological outcomes.

Research Procedure

The research procedure began with data collection, preparation, and integration from ecological databases, climate repositories, remote sensing sources, and biodiversity monitoring systems. Environmental datasets were standardized to ensure compatibility across different spatial and temporal scales (Redowan et al., 2025; Verma et al., 2025). The collected information was organized into a comprehensive database containing climate variables, ecosystem characteristics, biodiversity indicators, and ecosystem service measurements. Data quality assessment was conducted to identify missing information, inconsistencies, and potential sources of uncertainty before modeling analysis.

The second stage involved the development and calibration of the dynamic ecosystem service model. Ecological variables were incorporated into the modeling framework to represent interactions among climate conditions, biodiversity changes, ecosystem processes, and human influences. Climate change scenarios were simulated using projected environmental conditions to evaluate possible future ecosystem trajectories. Model parameters were adjusted through calibration procedures using historical ecological observations to improve prediction accuracy.

The third stage involved simulation analysis and evaluation of ecosystem responses under different climate change scenarios. The model generated projections regarding changes in ecosystem service availability, biodiversity vulnerability, habitat transformation, and ecological resilience. Comparative analysis was conducted among different scenarios to identify ecosystems at higher risk and determine potential conservation interventions.

Sensitivity analysis was performed to identify the most influential environmental factors affecting ecosystem stability and service provision.

The final stage involved interpretation of modeling results and translation of scientific findings into policy recommendations for biodiversity conservation. The research team analyzed model outputs to identify adaptive management strategies, conservation priorities, and climate resilience measures. The findings were evaluated through ecological and policy perspectives to ensure their relevance for decision-makers. The integration of dynamic modeling results with conservation planning provided a scientific foundation for developing proactive strategies to maintain biodiversity and ecosystem services under rapid climate change conditions.

Instruments, and Data Collection Techniques

The study utilized several scientific instruments and analytical tools to collect, process, and model ecosystem service dynamics. Climate datasets were obtained from global and regional climate databases, including historical temperature, precipitation, extreme weather events, and projected climate scenarios (Cohen et al., 2024). Remote sensing instruments and geographic information system (GIS) platforms were used to analyze land-use changes, habitat distribution, vegetation dynamics, and spatial patterns of ecosystem transformation. These instruments enabled the integration of environmental variables into the dynamic modeling framework.

Ecological assessment instruments were used to measure biodiversity conditions and ecosystem service performance. Biodiversity indicators included species richness, habitat suitability, population trends, and ecological connectivity metrics. Ecosystem service indicators measured carbon storage capacity, water regulation potential, soil protection functions, and habitat quality. The indicators were selected based on established ecological assessment frameworks to ensure consistency and scientific reliability in evaluating ecosystem responses under climate change scenarios.

Computational modeling instruments included ecological simulation software, statistical analysis platforms, and spatial modeling applications. The dynamic model was developed using integrated environmental datasets to simulate interactions among climate variables, biodiversity responses, ecosystem functions, and conservation strategies. Model validation instruments involved accuracy assessment, sensitivity analysis, and comparison between simulated outputs and observed ecological conditions. These procedures ensured that the model generated reliable predictions and meaningful interpretations for biodiversity conservation planning.

RESULTS AND DISCUSSION

The study analyzed integrated ecological datasets derived from climate records, biodiversity monitoring systems, remote sensing observations, and ecosystem service assessments. The dataset covered a 30-year historical period combined with future climate projections under different emission scenarios. The analysis incorporated 500 biodiversity indicators, including species distribution, habitat suitability, and population trends, alongside ecosystem service variables such as carbon sequestration, water regulation, soil conservation, and habitat provision. The spatial dataset represented forest ecosystems, freshwater systems, coastal environments, and agricultural landscapes exposed to different levels of climate vulnerability. The integration of multiple environmental datasets enabled the dynamic model to evaluate long-term ecosystem responses under changing climatic conditions.

Table 1. Descriptive Characteristics of Ecological Datasets Used in Dynamic Ecosystem Service Modeling

Data Components	Indicators	Dataset Coverage	Measurement Unit
Climate variables	Temperature, precipitation, extreme events	30-year historical and projected scenarios	°C, mm/year, frequency
Biodiversity indicators	Species richness, habitat suitability, population trends	500 species indicators	Index value
Land-use characteristics	Forest cover, agricultural expansion, habitat conversion	Multi-temporal satellite observations	km² and percentage
Ecosystem services	Carbon storage, water regulation, soil protection	Four ecosystem types	Service index
Conservation variables	Protected areas, management interventions	Selected conservation landscapes	Policy and ecological indicators

The descriptive analysis demonstrated significant variation in ecosystem responses across different climate scenarios. The baseline assessment showed that forest ecosystems provided the highest ecosystem service capacity, particularly in carbon storage and habitat protection, with an average ecosystem service index of 82.4 (SD = 6.7). Freshwater ecosystems showed moderate service capacity with an average index of 74.8 (SD = 8.1), while agricultural landscapes demonstrated greater sensitivity to climate variability with an average index of 68.5 (SD = 9.3). The results indicate that ecosystem resilience differs according to ecological characteristics, biodiversity complexity, and exposure to environmental pressures.

Table 2. Baseline Ecosystem Service Performance Across Different Ecosystem Types

Ecosystem Type	Carbon Storage Index	Water Regulation Index	Biodiversity Support Index	Overall Ecosystem Service Score
Forest ecosystems	86.7	81.5	79.0	82.4
Freshwater ecosystems	71.4	84.2	68.8	74.8
Coastal ecosystems	76.2	72.5	73.9	74.2
Agricultural landscapes	64.8	67.3	73.4	68.5

The descriptive findings reveal that ecosystem services respond differently to climate pressures depending on ecosystem structure, biodiversity composition, and environmental sensitivity. Forest ecosystems demonstrated relatively higher resilience because of their capacity to regulate climate processes, maintain habitat availability, and support diverse biological communities. Agricultural landscapes showed greater vulnerability because climate variability directly influenced productivity, soil conditions, and water availability. These patterns demonstrate that ecosystem service dynamics are strongly influenced by ecological complexity and adaptive capacity.

The scenario simulations indicated that rapid climate change may significantly alter ecosystem service availability over time. Under a high-emission scenario, ecosystem service scores declined across all ecosystem categories, with the greatest reduction occurring in biodiversity support functions. Forest ecosystems experienced an estimated 18.6% reduction in habitat suitability, while agricultural landscapes experienced a 25.4% decline in productivity-related ecosystem functions. The findings demonstrate that climate change creates uneven

ecological impacts, requiring conservation strategies tailored to specific ecosystem vulnerabilities.

The dynamic modeling analysis generated future projections of ecosystem service changes under three climate scenarios representing moderate, intermediate, and severe climate pressures. The simulation results showed that ecosystem degradation increased progressively as climate conditions intensified. Under the moderate scenario, ecosystem service capacity declined by approximately 8.7%, while intermediate and severe scenarios produced reductions of 15.9% and 24.6%, respectively. These results illustrate the cumulative effects of climate change on ecological functions and highlight the importance of early conservation interventions.

The biodiversity assessment revealed substantial changes in species distribution patterns under future climate scenarios. Approximately 23% of analyzed species indicators showed potential habitat contraction under severe climate conditions, while 17% demonstrated shifts toward more suitable ecological zones. Species with narrow habitat requirements experienced the highest vulnerability, particularly within fragmented ecosystems. The modeling results indicate that biodiversity responses are closely connected with ecosystem service stability and environmental adaptability.




Figure 1. Overall Ecosystem Service Capacity Loss

Inferential analysis was conducted to evaluate relationships between climate variables, biodiversity responses, and ecosystem service changes. Regression analysis demonstrated that temperature increase and precipitation variability significantly influenced ecosystem service performance. Temperature change showed a negative relationship with ecosystem stability ($\beta = -0.52$, $p < 0.001$), while precipitation variability also significantly affected ecosystem resilience ($\beta = -0.37$, $p < 0.001$). The statistical model explained 63.8% of the variation in ecosystem service changes, indicating that climate-related variables represent major determinants of ecological transformation.

Sensitivity analysis confirmed that biodiversity indicators and habitat connectivity significantly influenced ecosystem resilience under changing climate conditions. Biodiversity richness positively predicted ecosystem service stability ($\beta = 0.46$, $p < 0.001$), while habitat fragmentation negatively influenced ecological performance ($\beta = -0.41$, $p < 0.001$). The findings demonstrate that maintaining biodiversity and ecological connectivity can reduce climate-related risks and improve ecosystem adaptation capacity.

The relationship analysis demonstrated interconnected effects among climate change, biodiversity conditions, and ecosystem service performance. Climate pressures influenced biodiversity patterns, which subsequently affected ecosystem functions and service availability.

The structural relationship model indicated that climate change had both direct and indirect effects on ecosystem services through biodiversity alteration. The results emphasize that biodiversity conservation is not only a protection strategy for species but also an essential mechanism for maintaining ecosystem functionality.

The analysis further revealed that conservation interventions strengthened ecosystem resilience by reducing negative climate impacts. Protected areas, habitat restoration programs, and adaptive land management practices demonstrated positive relationships with ecosystem stability. Conservation intensity showed a significant positive association with ecosystem service maintenance ($r = 0.67$, $p < 0.001$). These findings indicate that policy interventions can modify ecological trajectories and improve ecosystem responses under future climate scenarios.

A detailed case study was conducted on a tropical forest ecosystem experiencing increasing climate pressure and biodiversity decline. The selected ecosystem represented a high-value conservation landscape due to its role in carbon storage, habitat protection, and regional climate regulation. Historical data indicated that the ecosystem experienced changes in vegetation structure, species distribution, and ecosystem service capacity due to temperature increases, rainfall variability, and land-use transformation. The dynamic model was applied to simulate future ecological conditions and evaluate potential conservation strategies.

The case study simulation demonstrated that unmanaged climate impacts could significantly reduce ecosystem functionality over the next several decades. Under the severe climate scenario, the forest ecosystem showed a projected decline of 21.3% in biodiversity support capacity and a 17.8% reduction in carbon storage potential. Conservation intervention scenarios, including habitat restoration and protected area expansion, reduced these losses by approximately 12.5%. The results demonstrate the importance of adaptive conservation planning based on predictive ecological modeling.

The case study findings explain that ecosystem resilience depends strongly on the interaction between environmental pressures and management capacity. The tropical forest ecosystem maintained higher ecological stability when conservation interventions reduced habitat fragmentation and supported species connectivity. Dynamic modeling revealed that proactive management strategies could significantly alter future ecological outcomes by reducing vulnerability and enhancing ecosystem recovery potential.

The findings also demonstrate the practical value of predictive modeling for conservation decision-making. Traditional conservation approaches often respond after ecological degradation occurs, whereas dynamic models enable policymakers to anticipate future risks and evaluate alternative strategies. The case study illustrates that integrating climate projections with biodiversity assessments provides more effective guidance for long-term ecosystem management.

The overall results demonstrate that dynamic modeling provides valuable insights into how ecosystem services respond to rapid climate change scenarios. The findings indicate that climate pressures, biodiversity changes, and ecosystem functions are strongly interconnected, requiring integrated approaches for effective conservation planning. Ecosystem vulnerability varies across landscapes, emphasizing the need for targeted policies based on ecological characteristics and future risk projections.

The study highlights that biodiversity conservation strategies should incorporate predictive modeling approaches to support adaptive environmental governance. Dynamic ecosystem service models allow decision-makers to evaluate potential consequences of climate change and identify interventions capable of maintaining ecological resilience. The findings provide evidence that science-based modeling can strengthen biodiversity conservation policies by connecting ecological forecasting with practical management decisions (Elleaume et al., 2025; Grajales Noreña et al., 2024).

The findings of this study demonstrate that dynamic modeling provides a comprehensive approach for understanding ecosystem service responses under rapidly changing climate conditions. The results reveal that climate variables, biodiversity patterns, habitat conditions, and ecosystem functions are strongly interconnected, creating complex ecological responses that cannot be adequately explained through static assessment methods. The modeling simulations indicate that increasing temperature, precipitation instability, and environmental disturbances contribute to significant reductions in ecosystem service capacity, particularly in biodiversity support, habitat stability, and ecological productivity. These findings confirm that climate change represents not only an environmental challenge but also a systemic disruption affecting the ability of ecosystems to sustain essential services.

The analysis demonstrates that ecosystem responses to climate change vary significantly depending on ecological characteristics and adaptive capacity. Forest ecosystems showed relatively higher resilience due to their complex biological structures, carbon storage capacity, and habitat connectivity, whereas agricultural landscapes experienced greater vulnerability because of their dependence on climatic stability and resource availability (Hulme et al., 2025; Pyron et al., 2024). The results indicate that ecosystem resilience is determined by the interaction between biodiversity complexity, environmental conditions, and management practices. These findings emphasize that conservation strategies should avoid generalized approaches and instead consider ecosystem-specific vulnerabilities and adaptive mechanisms.

The study further reveals that biodiversity conservation plays a central role in maintaining ecosystem service stability under future climate scenarios. The inferential analysis showed that biodiversity richness and habitat connectivity significantly influenced ecological resilience, suggesting that species conservation contributes directly to maintaining ecosystem functions. The modeling results indicate that ecosystems with higher biodiversity levels possess greater capacity to absorb environmental disturbances and recover from climate-related pressures. This finding reinforces the importance of integrating biodiversity protection into broader climate adaptation strategies.

The case study findings provide practical evidence that predictive modeling can support more effective conservation decision-making. The simulation of intervention scenarios demonstrated that proactive strategies, including habitat restoration, protected area expansion, and adaptive ecosystem management, could reduce projected ecological losses. The results indicate that dynamic models provide policymakers with valuable information regarding possible future conditions and potential management outcomes. The study confirms that integrating ecological forecasting with conservation planning can improve the effectiveness of biodiversity protection efforts under uncertain climate futures.

The findings are consistent with previous ecological studies demonstrating that climate change significantly affects biodiversity patterns, ecosystem functions, and the provision of ecosystem services. Existing research has established that rising temperatures, altered precipitation patterns, and extreme climatic events contribute to habitat degradation and species displacement. The present study supports these conclusions while extending previous knowledge by demonstrating how dynamic modeling can capture interactions among multiple ecological components over time. The findings provide additional evidence that climate impacts should be understood through integrated system perspectives rather than isolated environmental indicators.

Previous research on ecosystem services has frequently focused on measuring current ecological conditions or estimating the economic value of environmental benefits. These approaches have contributed significantly to understanding the importance of ecosystems for human well-being but often provide limited insights into future ecological trajectories (Hulme et al., 2025; Pyron et al., 2024). The current study differs by applying dynamic simulations to evaluate how ecosystem services may change under different climate scenarios. This approach

enables researchers and policymakers to move from descriptive assessments toward predictive analyses capable of informing long-term conservation planning.

Studies involving ecological modeling have introduced various predictive approaches, including species distribution models, climate impact models, and landscape simulations. These methods have successfully predicted specific ecological responses but often emphasize individual components such as species distribution or habitat suitability (Liu, Wang, et al., 2024; Umwali et al., 2025). The present research contributes by integrating biodiversity indicators, ecosystem service functions, climate projections, and conservation interventions within a unified framework. The findings demonstrate that comprehensive models can provide more realistic representations of ecological systems by incorporating multiple interacting factors.

The study also differs from previous research by emphasizing the policy relevance of dynamic ecological modeling. Many ecological models generate scientific predictions but provide limited translation into practical conservation strategies. The current findings demonstrate that modeling outcomes can guide decision-making by identifying vulnerable ecosystems, evaluating management alternatives, and predicting conservation outcomes. This contribution highlights the importance of linking ecological science with environmental governance to address complex challenges associated with rapid climate change.

The findings signify a fundamental shift in understanding ecosystem management from reactive conservation toward predictive and adaptive approaches. Traditional conservation strategies often respond to ecological degradation after significant changes have already occurred. The results of this study demonstrate that dynamic modeling enables early identification of potential risks and provides opportunities to implement preventive measures before irreversible ecological damage occurs. This represents an important transformation in biodiversity conservation by emphasizing anticipation, flexibility, and evidence-based planning.




Figure 2. Evolution of Ecosystem Management: From Reactive to Predictive & Adaptive Approaches

The results also signify that ecosystem services should be understood as dynamic processes rather than fixed ecological benefits. Climate change continuously modifies ecological relationships, affecting species interactions, habitat conditions, and ecosystem functions. The findings indicate that ecosystem services are shaped by ongoing interactions between environmental pressures and biological responses. This perspective challenges static approaches and highlights the importance of considering temporal changes when evaluating ecological sustainability.

The study further signifies that biodiversity conservation is directly connected to climate resilience and ecosystem stability. Biodiversity is not merely a conservation objective focused on protecting individual species but also a critical mechanism supporting ecosystem functionality. The findings demonstrate that maintaining diverse biological communities strengthens ecosystem capacity to adapt to environmental disturbances. This interpretation suggests that conservation policies should prioritize biodiversity as a foundation for maintaining ecosystem services under future climate scenarios.

The research findings signify the growing importance of integrating scientific modeling into environmental governance. Dynamic models provide a bridge between ecological knowledge and policy decisions by translating complex environmental processes into actionable information. The results indicate that effective conservation requires collaboration among scientists, policymakers, and resource managers. This integration represents a broader transformation toward adaptive governance capable of responding to rapidly changing ecological conditions.

The practical implication of this study is that policymakers should incorporate dynamic modeling approaches into biodiversity conservation planning. Traditional conservation decisions based solely on historical ecological conditions may become insufficient under rapidly changing climate conditions. The findings suggest that predictive models can assist governments and conservation organizations in identifying future ecological risks, prioritizing vulnerable ecosystems, and allocating resources more effectively. Dynamic modeling therefore provides a scientific foundation for developing forward-looking environmental policies.

The findings also have implications for ecosystem management practices. Conservation programs should move beyond species-focused protection toward integrated ecosystem-based approaches that consider climate adaptation, habitat connectivity, and ecosystem service maintenance. The results demonstrate that protecting biodiversity contributes to broader ecological resilience by maintaining essential ecosystem functions. Management strategies should therefore incorporate climate projections and ecological simulations when designing restoration programs and protected area policies.

The study provides implications for environmental governance by demonstrating the importance of evidence-based decision-making. Policymakers frequently face uncertainty when responding to climate change because ecological systems involve complex interactions and unpredictable outcomes. Dynamic modeling reduces this uncertainty by allowing decision-makers to evaluate possible scenarios before implementing interventions. The findings suggest that integrating scientific forecasting into policy development can improve the effectiveness and sustainability of conservation strategies.

The broader implication concerns the future direction of climate adaptation and biodiversity conservation. Ecosystem management must increasingly consider long-term ecological transformations rather than short-term environmental changes. The results indicate that maintaining ecosystem services under climate change requires continuous monitoring, adaptive policies, and interdisciplinary collaboration. The integration of modeling, ecological science, and governance frameworks can support more resilient ecosystems and sustainable human-environment relationships.

The observed ecosystem changes can be explained by the complex interactions between climate variables and ecological processes. Temperature increases and precipitation instability directly influence species survival, habitat conditions, and ecosystem productivity. Climate change alters environmental conditions faster than many ecosystems can naturally adapt, resulting in reduced ecological stability. The findings demonstrate that climate variables represent fundamental drivers shaping ecosystem service dynamics and biodiversity outcomes.

The differences in ecosystem vulnerability can be explained by variations in ecological complexity and adaptive capacity. Forest ecosystems generally possess greater resilience because they contain diverse species interactions, complex habitat structures, and stronger

regulatory functions (Aghaloo & Sharifi, 2025; Rasmus et al., 2024). Agricultural landscapes are more vulnerable because they often depend on simplified ecological systems and are directly influenced by climatic fluctuations. The results reflect the importance of ecosystem structure in determining responses to environmental stress.

The positive influence of biodiversity on ecosystem resilience can be explained through ecological complementarity and functional diversity mechanisms. Diverse biological communities provide multiple ecological functions that increase system stability under changing conditions. Species interactions contribute to resource regulation, nutrient cycling, and habitat maintenance. The findings demonstrate that biodiversity loss can create cascading effects that reduce ecosystem capacity to provide essential services.

The effectiveness of dynamic modeling can be explained by its ability to represent ecological complexity through simulation and scenario analysis. Ecosystems involve multiple interacting variables that change over time, making direct prediction difficult through conventional methods. Dynamic models integrate different environmental factors and simulate possible future conditions, allowing researchers to understand potential ecological trajectories. The findings demonstrate that modeling approaches are valuable because they capture both current conditions and future uncertainties.

Future research should expand dynamic ecosystem modeling by incorporating broader spatial scales, longer temporal periods, and more diverse ecological systems. The current study provides valuable insights into ecosystem responses under climate scenarios, but additional research is needed to examine variations across different geographical regions and ecological contexts. Comparative studies involving multiple ecosystems could improve understanding of how different environmental systems adapt to climate pressures.

Future investigations should integrate advanced technologies such as artificial intelligence, machine learning, and high-resolution remote sensing into ecosystem service modeling. These technologies may improve prediction accuracy by processing larger ecological datasets and identifying complex patterns within environmental changes. Research should focus on developing more adaptive models capable of incorporating real-time ecological information and improving conservation forecasting.

Further studies should examine the relationship between dynamic modeling outputs and actual policy implementation outcomes. Research is needed to evaluate whether predictive ecological information effectively improves conservation decisions and management effectiveness. Understanding how policymakers interpret and apply modeling results will strengthen the connection between scientific research and environmental governance.

The future of biodiversity conservation requires continuous integration between ecological modeling, technological innovation, and adaptive policy frameworks. Climate change will continue to create new ecological challenges that require flexible and science-based responses. The findings of this study suggest that dynamic modeling represents an important foundation for future conservation strategies by enabling proactive management, reducing uncertainty, and supporting long-term ecosystem resilience.

CONCLUSION

The most important finding of this study is that dynamic modeling provides a powerful approach for understanding the long-term interactions between climate change, biodiversity dynamics, and ecosystem service sustainability. The distinctive contribution of the findings lies in demonstrating that ecosystem responses cannot be accurately understood through static assessments because ecological systems continuously evolve through complex interactions among climate variables, biological processes, and human interventions. The research reveals that biodiversity conservation is not only essential for protecting species diversity but also functions as a critical mechanism for maintaining ecosystem resilience and service provision

under future climate pressures. The simulation results demonstrate that proactive conservation strategies, including habitat restoration, ecological connectivity improvement, and adaptive management, can significantly reduce potential ecosystem degradation. These findings provide important evidence that predictive ecological modeling can transform conservation planning from reactive responses toward anticipatory decision-making.

The major contribution of this research is primarily methodological through the development of an integrated dynamic modeling framework that combines climate scenario analysis, biodiversity assessment, ecosystem service evaluation, and conservation policy considerations. The study extends existing ecological modeling approaches by connecting environmental forecasting with practical decision-making processes, allowing policymakers to evaluate potential conservation outcomes before implementing interventions. The framework provides a comprehensive approach for analyzing ecosystem trajectories by incorporating multiple ecological variables rather than focusing on isolated indicators. The research contributes to environmental science by demonstrating how computational modeling can support adaptive governance, improve biodiversity conservation strategies, and strengthen evidence-based policy development under uncertain climate conditions.

The study acknowledges several limitations that provide directions for future research. The modeling framework depends on the availability, accuracy, and spatial resolution of ecological datasets, which may influence the reliability of long-term projections. The research primarily focused on selected ecosystem types and climate scenarios, limiting the generalizability of findings across all ecological regions and environmental contexts. Future studies should incorporate larger geographical areas, higher-resolution environmental data, and more complex ecological interactions to improve predictive accuracy. Further research should integrate artificial intelligence, machine learning techniques, and real-time ecological monitoring systems to enhance dynamic modeling capabilities. Longitudinal studies are also needed to evaluate how predicted ecosystem changes correspond with actual ecological responses and how modeling outputs can be effectively incorporated into global biodiversity conservation policies.

DECLARATION OF AI AND AI ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

During the preparation of this manuscript, the author(s) utilized Google Gemini solely for language translation and linguistic refinement purposes. All outputs generated by the tool were thoroughly reviewed, edited, and verified by the author(s) to ensure accuracy, clarity, and alignment with the original intent. The author(s) accept full responsibility for the integrity and content of the final publication.

AUTHOR CONTRIBUTIONS

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; In-vestigation.

Author 3: Data curation; Investigation.

Author 4: Formal analysis; Methodology; Writing - original draft.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

Abedin, I., Mukherjee, T., Kim, A. R., Kim, H.-W., Lee, S. R., & Kundu, S. (2025). Shifting shadows: Assessing the habitat and climate change response of binturong (Arctictis binturong) in the conservation landscape of the Asian continent. Ecological Informatics, 85, 102941. https://doi.org/https://doi.org/10.1016/j.ecoinf.2024.102941

Aghaloo, K., & Sharifi, A. (2025). Balancing priorities for a sustainable future in cities: Land use change and urban ecosystem service dynamics. Journal of Environmental Management, 382, 125460. https://doi.org/https://doi.org/10.1016/j.jenvman.2025.125460

Bai, J., Wang, X., Zhou, C., Zhou, J., Tu, Y., & Sun, Z. (2025). Integrative analysis of ecosystem health, ecosystem services and species habitat suitability can enhance the spatial configuration of protected areas. Ecological Indicators, 181, 114457. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114457

Banerjee, O., Cicowiez, M., Honeck, E. C., Muthukumara, M. S., & Stapleton, K. A. (2025). Investing in forests Enhances ecosystem services and economic Growth in Cambodia: Evidence from the Integrated Economic-Environmental Modelling (IEEM) approach. Ecosystem Services, 71, 101695. https://doi.org/https://doi.org/10.1016/j.ecoser.2024.101695

Bazzato, E., Calvia, G., Marignani, M., Ruggero, A., & Lozano, V. (2024). Senecio inaequidens DC. will thrive in future climate: A case study in a Mediterranean biodiversity hotspot. Ecological Informatics, 82, 102783. https://doi.org/https://doi.org/10.1016/j.ecoinf.2024.102783

Bianconi, A., Vascon, S., Furlan, E., & Critto, A. (2025). Leveraging artificial intelligence methods to map seagrass ecosystems in Italian Seas: Tackling human impact and climate change. Environmental Modelling & Software, 194, 106678. https://doi.org/https://doi.org/10.1016/j.envsoft.2025.106678

Bizama, G., Arismendi, I., Uribe-Rivera, D. E., Olivos, J. A., Jan, A., Valdovinos, C., & Urrutia, R. (2025). Climate change and freshwater biodiversity refuges in the Chilean Mediterranean Ecoregion. Science of The Total Environment, 1004, 180815. https://doi.org/https://doi.org/10.1016/j.scitotenv.2025.180815

Cohen, J. G., Enander, H. D., Bassett, T. J., Wilton, C. M., & Cole-Wick, A. A. (2024). Using GIS-based multicriteria decision analysis to prioritize invasive plant treatment: A creative solution for a pernicious problem. Ecological Modelling, 495, 110807. https://doi.org/https://doi.org/10.1016/j.ecolmodel.2024.110807

de Araujo Fonseca, F. N., & da Cunha Bustamante, M. M. (2025). The importance of indigenous territories for the provision of ecosystem services: A case study in the Brazilian Cerrado-Amazon Transition. Ecosystem Services, 72, 101706. https://doi.org/https://doi.org/10.1016/j.ecoser.2025.101706

Elleaume, N., Locatelli, B., Makowski, D., Vallet, A., Poulenard, J., Oszwald, J., & Lavorel, S. (2025). Uncertainties in future ecosystem services under land and climate scenarios: The case of erosion in the Alps. Ecological Modelling, 502, 111041. https://doi.org/https://doi.org/10.1016/j.ecolmodel.2025.111041

Ghasemi, K., Dolatkhahi, K., Qelichi, M. M., & Azadi, H. (2025). Assessing urban management resilience in response to rapid urbanization and climate change: A case study of Tehran. Urban Climate, 64, 102678. https://doi.org/https://doi.org/10.1016/j.uclim.2025.102678

González Hernández, M. M., León, C. J., García, C., & Lam-González, Y. E. (2023). Assessing the climate-related risk of marine biodiversity degradation for coastal and marine tourism. Ocean & Coastal Management, 232, 106436. https://doi.org/https://doi.org/10.1016/j.ocecoaman.2022.106436

Grajales Noreña, S., Bernal, G., Cardona, O. D., Rincón, D. F., & Carreño, M. L. (2024). Holistic evaluation of climate risk to prioritise adaptation measures for ecosystems. International Journal of Disaster Risk Reduction, 109, 104593. https://doi.org/https://doi.org/10.1016/j.ijdrr.2024.104593

Hulme, P. E., Lieurance, D., Richardson, D. M., & Robinson, T. B. (2025). Multiple targets of the Global Biodiversity Framework must be addressed to manage invasive alien species in protected areas. NeoBiota, 99, 149–170. https://doi.org/https://doi.org/10.3897/neobiota.99.152680

Kwak, Y., & Chen, S. (2025). Integrating seasonal climate variability and spatial accessibility in ecosystem service value assessment for optimized NbS allocation. Urban Climate, 59, 102314. https://doi.org/https://doi.org/10.1016/j.uclim.2025.102314

Lathouwers, M., Beenaerts, N., Evens, R., Artois, T., Conway, G., Henderson, I., Shewring, M., Cross, T., Ulenaers, E., & Dendoncker, N. (2025). Impact of future climate change and land-use change on habitat suitability for a long-distance avian migrant under diverse socioeconomic-emission scenarios. Global Ecology and Conservation, 62, e03777. https://doi.org/https://doi.org/10.1016/j.gecco.2025.e03777

Lin, F., Zhu, M., Dong, X., Ling, S., Luan, B., & Pan, G. (2025). Revealing future effectiveness of protected areas for biodiversity conservation in the Guangdong-Hong Kong-Macao Greater Bay Area. Landscape and Urban Planning, 255, 105260. https://doi.org/https://doi.org/10.1016/j.landurbplan.2024.105260

Lin, G.-Z., & Chiang, L.-C. (2025). Balancing ecosystem services and climate resilience: A case study of the Gaoping River Basin, Taiwan. Environmental and Sustainability Indicators, 27, 100745. https://doi.org/https://doi.org/10.1016/j.indic.2025.100745

Lin, G., Lan, S., Chi, B., Wu, K., Liu, Y., & Li, X. (2025). Synergistic optimization of ecological supply–demand: Integrating urban green space ecosystem services and dual ecological demands across multiple scenarios. Ecological Indicators, 178, 114133. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114133

Lin, Y.-P., Lin, P.-C., Wuryandani, S., Lin, C.-M., & Ros, G. H. (2025). Projecting food-energy-water sustainability through ecosystem service modeling under climate and land use change in a subtropical agricultural watershed. Agricultural Water Management, 318, 109737. https://doi.org/https://doi.org/10.1016/j.agwat.2025.109737

Lin, Z., Peng, S., Ding, X., Ma, D., Zhang, R., Lyu, M., Shi, S., Niu, F., & Pan, X. (2025). Unveiling ecosystem service dynamics under land use/cover changes in a plateau lake basin: a trajectory-scenario coupled approach. Ecological Indicators, 178, 114118. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114118

Liu, Y., Huang, X., & Liu, Y. (2024). Detection of long-term land use and ecosystem services dynamics in the Loess Hilly-Gully region based on artificial intelligence and multiple models. Journal of Cleaner Production, 447, 141560. https://doi.org/https://doi.org/10.1016/j.jclepro.2024.141560

Liu, Y., Ji, X., Xu, S., Li, L., Pan, Y., Wu, J., & Zhang, Y. (2025). Unraveling the influence of climate and land use change on ecological networks based on ecosystem services priority conservation areas. Global Ecology and Conservation, 64, e03923. https://doi.org/https://doi.org/10.1016/j.gecco.2025.e03923

Liu, Y., Wang, J., Wu, Z., Li, S., & Dai, W. (2024). Identification of biodiversity priority conservation areas in China by integrating genetic, species and ecosystem diversity. Biological Conservation, 300, 110854. https://doi.org/https://doi.org/10.1016/j.biocon.2024.110854

Pintado, K., & Knoke, T. (2025). Spatializing land-use scenarios to inform decision-making in an Ecuadorian mountain forest landscape. Journal of Environmental Management, 388, 125956. https://doi.org/https://doi.org/10.1016/j.jenvman.2025.125956

Pyron, R. A., Mooers, A. Ø., & Lo, N. Y. S. (2024). Nature's value and biodiversity ethics in a changing world: Insights from a special issue and questions for the future. Biological Conservation, 298, 110757. https://doi.org/https://doi.org/10.1016/j.biocon.2024.110757

Rasmus, S., Yletyinen, J., Sarkki, S., Landauer, M., Tuomi, M., Arneberg, M. K., Bjerke, J. W., Ehrich, D., Habeck, J. O., Horstkotte, T., Kivinen, S., Komu, T., Kumpula, T., Leppänen, L., Matthes, H., Rixen, C., Stark, S., Sun, N., Tømmervik, H., … Eronen, J. T. (2024). Policy documents considering biodiversity, land use, and climate in the European Arctic reveal visible, hidden, and imagined nexus approaches. One Earth, 7(2), 265–279. https://doi.org/https://doi.org/10.1016/j.oneear.2023.12.010

Redowan, M., Hewson, M., Abir, K. A. M., Hasan, F. E., Dey, B., & Kanan, A. H. (2025). Climate-induced shifts in the spatial distribution of invasive weed Sungrass (Imperata cylindrica) in the Asia-Pacific. Geomatica, 77(2), 100068. https://doi.org/https://doi.org/10.1016/j.geomat.2025.100068

Rieder, J., Jahnke, M., André, C., Christiansen, H., de Wit, P. R., Faust, E., Green, L., Jonsson, P. R., Laikre, L., Laugen, A. T., Rafajlović, M., Sandström, A., Tomasini, M., & Volckaert, F. A. M. (2025). Seascape genomics: Assisting marine biodiversity management by combining genetic knowledge with environmental and ecological information. Marine Policy, 182, 106867. https://doi.org/https://doi.org/10.1016/j.marpol.2025.106867

Strange, N., Ermgassen, S. zu, Marshall, E., Bull, J. W., & Jacobsen, J. B. (2024). Why it matters how biodiversity is measured in environmental valuation studies compared to conservation science. Biological Conservation, 292, 110546. https://doi.org/https://doi.org/10.1016/j.biocon.2024.110546

Tan, C., Yang, Q., Xu, B., Li, X., & Wu, X. (2025). Integrating ecological resilience into forest conservation planning to inform spatiotemporal prioritization. Biological Conservation, 312, 111491. https://doi.org/https://doi.org/10.1016/j.biocon.2025.111491

Umwali, E. D., Chen, X., Ma, X., Guo, Z., Mbigi, D., Zhang, Z., Umugwaneza, A., Gasirabo, A., & Umuhoza, J. (2025). Integrated SSP-RCP Scenarios for Modeling the Impacts of Climate Change and Land Use on Ecosystem Services in East Africa. Ecological Modelling, 504, 111092. https://doi.org/https://doi.org/10.1016/j.ecolmodel.2025.111092

Verma, D., Kumar, V., Shukla, A. K., & Pandey, R. (2025). Water provisioning services under changing land use and climate scenarios in the lower Shivaliks of Uttarakhand, India. Ecological Indicators, 180, 114334. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114334

Wang, F., Yuan, X., Sun, Y., & Liu, Y. (2024). Species distribution modeling based on MaxEnt to inform biodiversity conservation in the Central Urban Area of Chongqing Municipality. Ecological Indicators, 158, 111491. https://doi.org/https://doi.org/10.1016/j.ecolind.2023.111491

Wang, L., Zheng, H., Chen, Y., & Huang, B. (2024). Ecological redline policy strengthens sustainable development goals through the strict protection of multiple ecosystem services. Global Ecology and Conservation, 56, e03306. https://doi.org/https://doi.org/10.1016/j.gecco.2024.e03306

Wang, Y., & Liu, H. (2025). Integrating landscape ecological risk and ecosystem services for ecological zoning and management in the Wuling Mountain Area. Ecological Indicators, 179, 114179. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114179

Weng, A., Liu, Q., Lin, Y., Muhammad Nizamani, M., Wen, L., Zhou, Y., Wang, H., & Li, B. (2024). Dynamic conservation strategies for protected areas of Fujian Province: From integrated perspective of the adaptability of habitat and carbon storage to climate. Ecological Indicators, 168, 112773. https://doi.org/https://doi.org/10.1016/j.ecolind.2024.112773

Wu, S., Guo, Z., Askar, A., Li, X., Hu, Y., Li, H., & Saria, A. E. (2024). Dynamic land cover and ecosystem service changes in global coastal deltas under future climate scenarios. Ocean & Coastal Management, 258, 107384. https://doi.org/https://doi.org/10.1016/j.ocecoaman.2024.107384

Yang, S., Peng, S., Li, X., Wei, X., Pan, Y., & Jiao, Y. (2024). Spatial heterogeneity and interacting intensity of drivers for trade-offs and synergies between carbon sequestration and biodiversity. Global Ecology and Conservation, 56, e03256. https://doi.org/https://doi.org/10.1016/j.gecco.2024.e03256

Yang, W., Tanner, C. C., Holland, P., & Qu, Z. (2025). Dynamic economic valuation of coastal wetland restoration: A nature-based solution for climate and biodiversity. Environmental Challenges, 20, 101182. https://doi.org/https://doi.org/10.1016/j.envc.2025.101182

Yu, L., & Ma, Q. (2025). Synergies between bird habitat and recreational ecosystem service for urban development and biodiversity conservation in a coastal metropolitan region. Ecological Indicators, 181, 114408. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.114408

Zhi, L., Li, X., Bai, J., & Shao, D. (2025). Prioritizing multifunctional conservation zones with dominant function based on comprehensive hotspots and bundles of ecosystem service. Ecological Indicators, 171, 113122. https://doi.org/https://doi.org/10.1016/j.ecolind.2025.113122

Copyright Holder :
© Achmad Agus Salim et al. (2026).

First Publication Right :
© Research of Scientia Naturalis

This article is under:

